

BORELIC PAIRS FOR STRATIFIED ALGEBRAS

KEVIN COULEMBIER AND RUIBIN ZHANG

ABSTRACT. We determine all values of the parameters for which the cell modules form a standard system, for a class of cellular diagram algebras including partition, Brauer, walled Brauer, Temperley-Lieb and Jones algebras. For this, we develop and apply a general theory of algebras with Borelic pairs. The theory is also applied to give new uniform proofs of the cellular and quasi-hereditary properties of the diagram algebras and to construct quasi-hereditary 1-covers, in the sense of Rouquier, with exact Borel subalgebras, in the sense of König. Another application of the theory leads to a proof that Auslander-Dlab-Ringel algebras admit exact Borel subalgebras.

CONTENTS

1. Introduction	1
2. Preliminaries	6
Part I. General theory	11
3. Pre-Borelic pairs	11
4. Borelic pairs	15
5. Base stratified algebras	21
Part II. Examples	24
6. Auslander-Dlab-Ringel algebras	24
7. Example from Lie theory	26
8. Algebras in diagram and partition categories	27
Part III. Faithfulness of covers	36
9. Covers of the diagram algebras	36
Appendix A. Stratified algebras	45
References	47

1. INTRODUCTION

The theory of quasi-hereditary algebras was initiated by Cline, Parshall and Scott in [CPS1] and provided a unified framework for studying modular representation theory of semisimple algebraic groups and the BGG category $\mathcal{O}$. A central role in this theory is played by the *standard modules* of quasi-hereditary algebras. Cellular algebras were introduced by Graham

2010 *Mathematics Subject Classification.* 16G10, 17B10, 20C05, 81R05.

Key words and phrases. stratified and quasi-hereditary algebras, exact Borel subalgebras, cellular and standardly based algebras, standard systems, quasi-hereditary covers, diagram algebras, Schur algebras.

and Lehrer in [GL] and include many *diagram algebras* such as Brauer, Iwahori-Hecke and BMW algebras. These algebras are of central importance in representation theory and low dimensional topology. Cellular algebras have a class of natural modules, known as the *cell modules*. For many values of their parameters, Brauer and BMW algebras are also quasi-hereditary and then the standard modules coincide with the cell modules. This remains true for a range of cellular algebras.

In [KN, HN], it is proved that, in most cases, the cell modules of Iwahori-Hecke algebras of type A behave as the standard modules of some quasi-hereditary algebra, even though the Iwahori-Hecke algebra is itself not quasi-hereditary. This is formulated into the statement that “the cell modules form a *standard system*”, see [DR2], and is equivalent to the condition that the algebra admits a *Schur algebra*, in the sense of [HHKP, Ro]. The result in [KN] extends to Brauer algebras, by [HP], to partition and BMW algebras, by [HHKP], and to Iwahori-Hecke algebras of type B , by [Ro]. An important consequence is that multiplicities in cell filtrations are well-defined.

In the present paper, we determine when the cell modules form a standard system for a variety of diagram algebras: partition, Brauer, walled Brauer, Temperley-Lieb and Jones algebras. In the cases of the Brauer and partition algebras, this reproduces (and completes) the corresponding results of [HP, HHKP]. We will achieve this by developing a construction which is in part inspired by the theory of exact Borel subalgebras.

In [Kö], König introduced the notion of an *exact Borel subalgebra* of a quasi-hereditary algebra, inspired by the Borel subalgebra of a semisimple Lie algebra. When an exact Borel subalgebra exists, the standard modules are given explicitly as the modules induced from the simple modules of that subalgebra. Not every quasi-hereditary algebra admits an exact Borel subalgebra. However, it was proved in [KKO] that every quasi-hereditary algebra over an algebraically closed field is Morita equivalent to a (quasi-hereditary) algebra admitting an exact Borel subalgebra. Although the proof is constructive, it would not be trivial to apply it to construct the Morita equivalent algebra and its exact Borel subalgebra for a given quasi-hereditary algebra. In fact, few explicit examples of Morita equivalent algebras with exact Borel subalgebras for prevalent quasi-hereditary algebras are known, see *e.g.* [Kö, PSW].

We will define a generalisation of the concept of exact Borel subalgebras, *viz.* *Borelic pairs* (B, H) of arbitrary algebras A , and develop the theory of algebras with such pairs. We prove that key structural properties of A are determined by those of H . For instance, if H is semisimple then A is quasi-hereditary, and, roughly speaking, cellularity of H implies cellularity of A . Furthermore, in the latter case, the cell modules of A form a standard system if and only if this is the case for H . These and further applications are more precisely discussed in Section 1.3.


We use this general theory to study the diagram algebras mentioned above, in particular for (i) determining when the cell modules form standard systems and (ii) constructing Morita equivalent algebras with exact Borel subalgebras. For any of the diagram algebras A , we construct an algebra C , satisfying a double centraliser property with A . The algebra C admits a Borelic pair, which allows to prove that it is cellular, determine when the cell modules form a standard system, and find out when it is quasi-hereditary. In the latter case, C admits an exact Borel subalgebra. In most cases, C is Morita equivalent to A and in the remaining cases, it is a 1-faithful cover in the sense of [Ro]. In some cases, C constitutes a previously unknown Schur algebra. To the best of our knowledge, this is the first purely

diagrammatic description of a Schur algebra. Aside from the new results on the algebras A , our constructions also yields a new proof of their cellularity and quasi-heredity.

1.1. Standardly based and base stratified algebras. In [DuR], Du and Rui introduced a generalisation of cellular algebras, called *standardly based algebras*. We will prove that any algebra over an algebraically closed field can be given at least one standardly based structure. Standardly based algebras also come with cell modules. When the standardly based algebra is cellular, the two types of cell modules coincide.

In [HHKP], Hartman *et al.* defined the concept of *cellularly stratified algebras*, giving a powerful tool to determine when cell modules form a standard system. We introduce the corresponding weaker notion of *base stratified algebras*, which is a significant simplification that is better compatible with the notion of Borelic pairs, and actually suffices for studying when the cell modules of a cellular algebra form a standard system.

FIGURE 1. Links between structures



1.2. Overview of the structures on algebras. The different structures on algebras we study are summarised in Figure 1. The notion of *properly stratified algebras* was introduced in [DL, KM]. The notion of *(standardly) stratified algebras* was introduced in [CPS2, Chapter 2]. The notion of *strongly standardly stratified algebras* was studied in [ADL], although they were called “standardly stratified” in that paper, leading to inconsistency with [CPS2]. The term “strongly standardly stratified algebras” was coined in [Fr], which also introduced *exactly standardly stratified algebras*, as “weakly properly stratified algebras”.

An arrow means that one structure implies the other one. For arrows between different versions of standardly stratified algebras, the (proper) standard modules remain the same. The arrow from quasi-hereditary to base stratified only holds when the field is algebraically closed and in this case the standard modules are mapped to the cell modules. A general base stratified algebras has standard modules, cell modules and proper standard modules. The first type of modules are different from the other two, unless the algebra is quasi-hereditary.

1.3. Borelic pairs. We generalise the notion in [Kö] of exact Borel subalgebras from quasi-hereditary algebras to exactly standardly stratified algebras, where the generalisation to properly stratified algebras was already introduced by Klucznik and Mazorchuk in [KM].

We further generalise this theory by introducing the notion of *Borelic pairs* (B, H) of an arbitrary algebra A . We prove that, when A admits a Borelic pair (B, H) , it is standardly stratified. Moreover, if H is quasi-local, A is strongly standardly stratified. We also consider a stronger notion of *exact Borelic pairs*, leading to exactly standardly stratified algebras. Then we find that, if H is quasi-local (resp. semisimple) A is properly stratified (resp. quasi-hereditary) and in each case B is an exact Borel subalgebra. If H is standardly based, A is base stratified, so in particular standardly based. Furthermore, the cell modules of H form a standard system if and only the standard modules of A form a standard system.

1.4. Overview of main applications. The following will be proved in Theorem 6.3.1.

Theorem 1. *Over a perfect field $\mathbb{k}$, the Auslander-Dlab-Ringel algebra of a finite dimensional unital $\mathbb{k}$ -algebra R is quasi-hereditary with exact Borel subalgebra.*

The weak assumption that $\mathbb{k}$ is perfect is not required for the quasi-heredity of ADR algebras, see [DR1], but our construction of the exact Borel subalgebra fails without it. This exact Borel subalgebra and the ones in Theorem 2 seem unrelated to the construction in [KKO], see e.g. Remark 8.4.8. The quasi-heredity of ADR algebras has the following consequence, see Theorem 6.4.1.

Corollary 1. *Any algebra over an algebraically closed field has a standardly based structure.*

The following theorem, which uses a total quasi-order $\preceq$ and a partial order $\leq$ on the set of simple modules, will be proved in Theorems 8.4.1, 8.5.1 and 9.2.1 and Corollary 8.6.3.

Theorem 2. *Consider an arbitrary field $\mathbb{k}$ and a fixed $\delta \in \mathbb{k}$. Let A be the partition algebra $\mathcal{P}_n(\delta)$, the Brauer algebra $\mathcal{B}_n(\delta)$, the walled Brauer algebra $\mathcal{B}_{r,s}(\delta)$, the Jones algebra $J_n(\delta)$ or the Temperley-Lieb algebra $\text{TL}_n(\delta)$, with $n, r, s \in \mathbb{N}$, $n > 1$, $r \geq 1$ and $s \geq 1$. Then*

- (1) A is cellular if $\mathcal{C}_A^3$;
- (2) $(A, \preceq)$ is exactly standardly stratified^(*) if $\mathcal{C}_A^1$;
- (3) $(A, \leq)$ is quasi-hereditary if $\mathcal{C}_A^1$ and $\mathcal{C}_A^2$;
- (4) A admits a cover C which is:
 - Morita equivalent to A if $\mathcal{C}_A^1$;
 - quasi-hereditary with exact Borel subalgebra if $\mathcal{C}_A^2$;
 - always exactly standardly stratified^(*) with exact Borel subalgebra;
 - cellular and base stratified if $\mathcal{C}_A^3$.

algebra A	$\mathcal{C}_A^1$	$\mathcal{C}_A^2$	$\mathcal{C}_A^3$
$\mathcal{P}_n(\delta)$	$\delta \neq 0$	$\text{char}(\mathbb{k}) \notin [2, n]$	$\emptyset$
$\mathcal{B}_n(\delta)$	$\delta \neq 0$ or n odd	$\text{char}(\mathbb{k}) \notin [2, n]$	$\emptyset$
$J_n(\delta)$	$\delta \neq 0$ or n odd	n odd and $\text{char}(\mathbb{k}) \notin [3, n]$ or n even and $\text{char}(\mathbb{k}) \notin \{2\} \cup [3, n/2]$	$x^i - 1$ splits over $\mathbb{k}$, for $i \in \{n, n-2, \dots\}$
$\text{TL}_n(\delta)$	$\delta \neq 0$ or n odd	$\emptyset$	$\emptyset$
$\mathcal{B}_{r,s}(\delta)$	$\delta \neq 0$ or $r \neq s$	$\text{char}(\mathbb{k}) \notin [2, \max(r, s)]$	$\emptyset$

When $\mathcal{C}_A^1$ is satisfied, but not $\mathcal{C}_A^2$, the algebra A is not quasi-hereditary, for any order.

(*) In case $A = J_n(\delta)$, even properly stratified.

When $\mathcal{C}_A^1$ is not satisfied, it follows from [KX2] that A is not quasi-hereditary, for any order. Hence, we obtained a Morita equivalent algebra with exact Borel subalgebra for all cases when A is quasi-hereditary. The quasi-heredity and cellularity of A in Theorem 2 were previously obtained by a variety of methods in [AST, CDDM, GL, KX2, Ma2, Xi]. Here we find a new unified proof, which also constructs the exact Borel subalgebras. The quasi-heredity of $J_n(\delta)$ has previously only been stated, in [KX2, Proposition 4.2], without the explicit conditions. Next we determine when the cell modules of A form a standard system.

Theorem 3. *If the field $\mathbb{k}$ is algebraically closed, all algebras A in Theorem 2 are cellular. The cell modules form a standard system if and only if the following condition is satisfied:*

algebra A	condition
$\mathcal{P}_n(\delta)$	$\delta \neq 0$ if $n = 2$; and $\begin{cases} \text{char}(\mathbb{k}) \notin \{2, 3\} & \text{or} \\ \text{char}(\mathbb{k}) = 3 \text{ and } n = 2 \end{cases}$
$\mathcal{B}_n(\delta)$	$\delta \neq 0$ if $n \in \{2, 4\}$; and $\begin{cases} \text{char}(\mathbb{k}) \notin \{2, 3\} & \text{or} \\ \text{char}(\mathbb{k}) = 3 \text{ and } n = 2 \end{cases}$
$J_n(\delta)$	$\delta \neq 0$ if $n \in \{2, 4\}$; and $\begin{cases} \text{char}(\mathbb{k}) \notin [3, n] & \text{if } n \text{ is odd} \\ \text{char}(\mathbb{k}) \notin \{2\} \cup [3, n/2] & \text{if } n \text{ is even} \end{cases}$
$\text{TL}_n(\delta)$	$\delta \neq 0$ if $n \in \{2, 4\}$
$\mathcal{B}_{r,s}(\delta)$	$\delta \neq 0$ if $(r, s) \in \{(1, 1), (2, 2)\}$; and $\begin{cases} \text{char}(\mathbb{k}) \notin \{2, 3\} & \text{or} \\ \text{char}(\mathbb{k}) = 3 \text{ and } \max(r, s) \in \{1, 2\} \end{cases}$

This follows from Theorems 8.4.3, 8.5.1 and 9.3.1. The results for $\mathcal{B}_n(\delta)$ were previously obtained in [HP] and most of the claim for $\mathcal{P}_n(\delta)$ was proved in [HHKP]. This theorem, together with [DR2, Theorem 2] and Lemma 5.3.1, yields the following consequence.

Corollary 2. *The cellular algebra A , under the condition in Theorem 3, admits a Schur algebra in the sense of [HHKP, Definition 12.1] or Definition 2.9.4. Moreover, the cell multiplicities in modules which admit cell filtrations are independent of the specific filtration.*

When the condition $\mathcal{C}_A^1$ in Theorem 2 is not satisfied and the cover C is hence not Morita equivalent to A , in most cases it is still a 1-faithful cover, thus a quasi-hereditary 1-cover in the sense of [Ro] when C is quasi-hereditary (*i.e.* when $\mathcal{C}_A^2$ is satisfied).

Theorem 4. *Consider an arbitrary field $\mathbb{k}$ and the cover C of A in Theorem 2, then C is a 1-faithful quasi-hereditary cover if $\mathcal{C}_A^2$ and the condition in the table is satisfied.*

algebra A	condition (along with $\mathcal{C}_A^2$)
$\mathcal{P}_n(\delta)$	$n \neq 2$ or $\delta \neq 0$
$\mathcal{B}_n(\delta)$	$n \notin \{2, 4\}$ or $\delta \neq 0$
$J_n(\delta)$	$n \notin \{2, 4\}$ or $\delta \neq 0$
$\text{TL}_n(\delta)$	$n \notin \{2, 4\}$ or $\delta \neq 0$
$\mathcal{B}_{r,s}(\delta)$	$(r, s) \notin \{(1, 1), (2, 2)\}$ or $\delta \neq 0$

Under these conditions, the algebra C is the Schur algebra predicted in Corollary 2.

This is proved in Theorems 9.2.2 and 8.5.1. If, for $A = \mathcal{B}_n(\delta)$, we take the stronger condition ‘ n even or $\delta \neq 0$ ’, but relax condition $\mathcal{C}_A^2$ to ‘ $\text{char}(\mathbb{k}) \notin \{2, 3\}$ ’, the Schur algebra of Corollary 2 is constructed in a more combinatorial way in [HK], see also [Bo] for the walled

Brauer algebra. Perhaps a combination of our methods with the ones in [HK] can be used to construct the Schur algebras of Corollary 2 in full generality.

Finally, we remark that our treatment of Theorems 1, 2 and 3 provide alternative proofs for several results in [AST, CDDM, DR1, GL, HHKP, HP, KX2, Ma2, Xi]. Our approach does not rely on any of the results in these papers.

2. PRELIMINARIES

We work over an arbitrary ground field $\mathbb{k}$. Unless specified otherwise, we make no assumptions on its characteristic and do not require it to be algebraically closed. A field is called *perfect* if every finite field extension of $\mathbb{k}$ is separable. Examples are algebraically closed fields, fields of characteristic zero and finite fields. We set $\mathbb{N} = \{0, 1, 2, \dots\}$.

2.1. Algebras and modules. By an algebra A over $\mathbb{k}$, we will always mean an *associative, unital, finite dimensional* algebra. All subalgebras of an algebra A are assumed to contain the identity 1_A . Modules over A are supposed to be unital, in the sense that 1_A acts as the identity. Furthermore, they are always assumed to be left modules and finitely generated, unless explicitly specified otherwise. The corresponding module category is denoted by $A\text{-mod}$. Two algebras A and B are *Morita equivalent* if there is an equivalence of categories

$$A\text{-mod} \cong B\text{-mod}.$$

We write this as $A \stackrel{M}{=} B$. A *local* algebra is an algebra with unique maximal left ideal. We say that an algebra is *quasi-local* if it is Morita equivalent to the Cartesian product of local algebras, so if there are no extensions between non-isomorphic simple modules.

2.2. Jacobson radical. The Jacobson radical $\text{rad}A$ of an algebra A is the intersection of all maximal left ideals, see [La, Chapter 4]. An algebra A is called *semisimple* if its Jacobson radical is zero, which is equivalent to the condition that A is the Cartesian product of simple algebras by [La, Theorem 4.14]. All finite dimensional unital algebras are *semi-primary*, meaning that $A/\text{rad}A$ is semisimple and $\text{rad}A$ is nilpotent, see [La, Theorem 4.15].

A $\mathbb{k}$ -algebra A is called *separable* if for any field extension $\mathbb{K}$ of $\mathbb{k}$, $A \otimes_{\mathbb{k}} \mathbb{K}$ is a semisimple $\mathbb{K}$ -algebra. Every simple algebra over a perfect field is separable. If A has a semisimple subalgebra S , such that $A = S \oplus \text{rad}A$ as vector spaces, we say that A is a *Wedderburn algebra*. Wedderburn's principal theorem states that any algebra such that $A/\text{rad}A$ is separable is Wedderburn, see [We, Exercise 9.3.1].

2.3. Idempotents. An *idempotent* $e \in A$ is *primitive* if the left A -module Ae is indecomposable. Two idempotents e and f in A are *equivalent* if and only if $Ae \cong Af$ as left A -modules, or equivalently if there are $a, b \in A$ such that $e = ab$ and $f = ba$.

For an algebra A , there is a one-to-one correspondence between the isomorphism classes of simple unital modules, the equivalence classes of primitive idempotents and the isomorphism classes of projective covers of the simple modules. So there is a (finite) set $\Lambda = \Lambda_A$ with corresponding simple module $L(\lambda)$, primitive idempotent e_λ and projective module $P(\lambda) := Ae_\lambda$ for each $\lambda \in \Lambda_A$, exhausting the classes irredundantly, such that

$$e_\lambda L(\lambda') \neq 0 \quad \Leftrightarrow \quad \lambda' = \lambda \quad \Leftrightarrow \quad \text{Hom}_A(P(\lambda), L(\lambda')) \neq 0.$$

2.4. Centraliser algebras. For an algebra C with idempotent e , the centraliser algebra of e is $C_0 = eCe$. We consider the pair of adjoint functors (G, F) :

$$(2.1) \quad \begin{array}{ccc} C\text{-mod} & \begin{array}{c} \xrightarrow{F=\text{Hom}_C(Ce, -)=e-} \\ \xleftarrow{G=Ce \otimes_{C_0} -} \end{array} & C_0\text{-mod}. \end{array}$$

Clearly we have $F \circ G \cong \text{Id}$ on $C_0\text{-mod}$, and F is exact while G is right exact. Sometimes it will be more convenient to consider $G' = \text{Hom}_{C_0}(eC, -)$, the right adjoint to F .

We briefly elaborate on three special cases of the above:

(a) We say that C is a *cover* of C_0 if the restriction of F to the category of projective C -modules is fully faithful. This is equivalent to demanding that the canonical morphism $C \rightarrow \text{End}_{C_0}(eC)^{\text{op}}$, or $C \rightarrow \text{End}_{C_0}(Ce)$ for the right C_0 -module Ce , is an isomorphism.

(b) Assume that the right C_0 -module Ce is projective and hence G an exact functor. Then we have an adjoint pair (G, F) of exact functors and consequently

$$(2.2) \quad \text{Ext}_C^k(G-, -) \cong \text{Ext}_{C_0}^k(-, F-), \quad \text{for all } k \geq 0.$$

(c) Assume that $C = CeC$, then every primitive idempotent of C is equivalent to one contained in eCe . Hence Ce is a projective generator for $C\text{-mod}$, so it follows that $G \circ F \cong \text{Id}$ on $C\text{-mod}$. Hence, in this case, C and C_0 are Morita equivalent.

2.5. Orders and partitions. A *partial quasi-order* $\preccurlyeq$ is a binary relation which is reflexive and transitive. When a partial quasi-order is also anti-symmetric, it is a *partial order*, and will usually be denoted by $\leq$. When $\preccurlyeq$ satisfies the condition that for any elements s, t in the set at least one of $s \preccurlyeq t$ or $t \preccurlyeq s$ holds, it is a *total quasi-order*, or simply a *quasi-order*.

For a partial quasi-order $\preccurlyeq$, we use the notation $s \prec t$ when $s \preccurlyeq t$ but $t \not\preccurlyeq s$, and $s \sim t$ when $s \preccurlyeq t$ and $t \preccurlyeq s$. An *extension* $\preccurlyeq_e$ of a partial quasi-order $\preccurlyeq$ is a partial quasi-order such that $s \preccurlyeq t$ implies $s \preccurlyeq_e t$, and such that $s \sim t$ if and only if $s \sim_e t$. In particular, an extension of a partial order remains a partial order.

An *n-decomposition* $\mathcal{Q}$ of a set S is an ordered disjoint union $S = \sqcup_{i=0}^n S_i$ into $n+1$ subsets. Decompositions are in natural bijection with total quasi-orders. For the decomposition $\mathcal{Q}$ the corresponding total quasi-order $\preccurlyeq_{\mathcal{Q}}$ on S is given by $s \preccurlyeq_{\mathcal{Q}} t$ if and only if $s \in S_i$ and $t \in S_j$ with $i \geq j$. To each decomposition $\mathcal{Q}$, we also associate a partial order $\leq_{\mathcal{Q}}$, defined by $s <_{\mathcal{Q}} t$ if and only if $s \in S_i$ and $t \in S_j$ with $i > j$. Note that we cannot obtain every partial order in this way.

We will use the term *partition* in two different situations. A *partition of the natural number* n is a (non-strictly) decreasing sequence of natural numbers adding up to n . When λ is a partition of n we write $\lambda \vdash n$. For a natural number p , $\lambda \vdash n$ is *p-regular* if the sequence of the partition does not contain p times the same non-zero number, see [Ja, Section 6.1]. For a field $\mathbb{k}$, we say that λ is $\mathbb{k}$ -regular if either $\text{char}(\mathbb{k}) = 0$ or λ is $\text{char}(\mathbb{k})$ -regular, and write this as $\lambda \vdash_{\mathbb{k}} n$. Secondly, we will use the notion of a *partition of a set*, which is a grouping of the elements into non-empty subsets (or equivalence classes).

2.6. Stratifying ideals. An idempotent ideal J in an algebra A is a two-sided ideal which satisfies $J^2 = J$. This means $J = AeA$ for some idempotent e , see *e.g.* [APT]. This idempotent e is not uniquely determined by J . However, it follows easily, see *e.g.* [APT], that $AeA = A\tilde{e}A$, for two idempotents e and $\tilde{e}$, implies

$$(2.3) \quad eAe \stackrel{M}{=} \tilde{e}A\tilde{e}.$$

2.6.1. We introduce some specific types of stratifying ideals, using in particular terminology from [CPS1, CPS2, Fr, KM]. An idempotent ideal $J = AeA$ in A is

- (0) an *exactly stratifying ideal* if:
 - the right A -module J_A is projective.
- (1) a *standardly stratifying ideal* if:
 - the A -module ${}_AJ$ is projective.
- (2) a *exactly standardly stratifying ideal* if:
 - the left A -module ${}_AJ$ and right A -module J_A are projective.
- (3) a *strongly standardly stratifying ideal* if:
 - the A -module ${}_AJ$ is projective and
 - the algebra eAe is quasi-local.
- (4) a *properly stratifying ideal* if:
 - the left A -module ${}_AJ$ and right A -module J_A are projective and
 - the algebra eAe is quasi-local.
- (5) a *heredity ideal* if:
 - the A -module ${}_AJ$ is projective and
 - the algebra eAe is semisimple.

All these ideals are stratifying in the sense of [CPS2, Definition 2.1.1]. A heredity ideal is properly stratifying by [APT, Corollary 5.3], the other relations in Figure 1 are by definition.

Remark 2.6.2. By equation (2.3), the definitions depend only on J and not on e .

2.6.3. We say that a chain of idempotent ideals

$$(2.4) \quad 0 = J_0 \subsetneq J_1 \subsetneq \cdots \subsetneq J_{m-1} \subsetneq J_m = A,$$

has length $m - 1$. With this convention, the trivial chain $0 = J_0 \subset J_1 = A$ has length 0. Chains of length k are in one-to-one correspondence with k -decompositions of Λ . The decomposition corresponding to a chain of idempotent ideals (2.4) is defined by setting $\lambda \in \Lambda_i$ for the minimal i for which $J_{i+1}L(\lambda) \neq 0$. Consequently, the chains of idempotent ideals of A are also in one-to-one correspondence with the total quasi-orders of Λ .

2.6.4. If for a chain (2.4), the ideal J_i/J_{i-1} is a standardly, exactly standardly, strongly standardly or properly stratifying ideal in A/J_{i-1} , the chain is also called standardly, exactly standardly, strongly standardly or properly stratifying. If each ideal J_i/J_{i-1} is a heredity ideal, the chain is called heredity.

We will take the convention of denoting the image of an idempotent $e \in A$, in the quotient A/J , for some idempotent ideal J again by e .

Remark 2.6.5. For a chain (2.4), we can choose idempotents f_i for $i \in \{1, \dots, m\}$ such that $J_i = Af_iA$, with $f_i f_j = f_i = f_j f_i$ if $i \leq j$ and $f_m = 1$. We define the algebras

$$A^{(j)} = f_{j+1}(A/J_j)f_{j+1}, \quad \text{for } j \in \{0, 1, \dots, m-1\},$$

which are only uniquely associated to the chain up to Morita equivalence, by (2.3).

2.7. Standardly stratified algebras. An algebra A with some partial quasi-order $\preccurlyeq$ on Λ will be denoted as $(A, \preccurlyeq)$.

Definition 2.7.1. Consider an algebra $(A, \preccurlyeq)$, for $\preccurlyeq$ a total quasi-order, and the chain (2.4) of idempotent ideals corresponding to $\preccurlyeq$. We say that $(A, \preccurlyeq)$ is

- (1) *standardly stratified*, if the chain (2.4) is standardly stratifying;

- (2) *exactly standardly stratified*, if the chain (2.4) is exactly standardly stratifying;
- (3) *strongly standardly stratified*, if $\preceq$ is a total order and the chain (2.4) is strongly standardly stratifying;
- (4) *properly stratified*, if $\preceq$ is a total order and the chain (2.4) is properly stratifying;
- (5) *quasi-hereditary*, if $\preceq$ is a total order and the chain (2.4) is heredity.

Remark 2.7.2. The trivial chain $0 = J_0 \subset J_1 = A$ is an exactly standardly stratifying chain for any algebra A . It is thus essential to specify the quasi-order $\preceq$ when speaking about types of standardly stratified algebras.

Remark 2.7.3. If $\preceq$ is a total order and the chain (2.4) corresponding to $\preceq$ is standardly (resp. exactly standardly) stratifying, it is automatically strongly standardly (resp. properly) stratifying, as the algebra $A^{(i)}$ will only have one simple module up to isomorphism.

There is also a module theoretic approach to standardly stratified algebras, where we no longer demand the orders to be total.

Definition 2.7.4. Consider an algebra $(A, \preceq)$ for some partial quasi-order $\preceq$ on Λ . For any $\lambda \in \Lambda$, let $L(\lambda)$ denote the corresponding simple A -module and $P(\lambda)$ its projective cover.

- (1) If there is a set of A -modules $\{S(\lambda), \lambda \in \Lambda\}$ such that for any $\lambda, \mu \in \Lambda$:
 - we have $[S(\lambda) : L(\mu)] = 0$ unless $\mu \preceq \lambda$, and
 - there is a surjection $P(\lambda) \twoheadrightarrow S(\lambda)$ such that the kernel has a filtration where the sections are isomorphic to modules $S(\nu)$ for $\lambda \prec \nu$,
 we say that $(A, \preceq)$ is *standardly stratified*.
- (2) If $(A, \preceq)$ is standardly stratified and there are modules $\{\overline{S}(\lambda), \lambda \in \Lambda\}$ with $[\overline{S}(\lambda) : L(\lambda)] = 1$ and $[\overline{S}(\lambda) : L(\mu)] = 0$ unless $\mu = \lambda$ or $\mu \prec \lambda$, such that each module $S(\lambda)$ has a filtration where the sections are isomorphic to modules $\overline{S}(\mu)$ with $\mu \sim \lambda$, we say that $(A, \preceq)$ is *exactly standardly stratified*.
- (3) If $\preceq$ is a partial order and $(A, \preceq)$ is standardly stratified as in (1), we say that $(A, \preceq)$ is *strongly standardly stratified*.
- (4) If $\preceq$ is a partial order and $(A, \preceq)$ is exactly standardly stratified as in (2), we say that $(A, \preceq)$ is *properly stratified*.
- (5) If $\preceq$ is a partial order, $(A, \preceq)$ is standardly stratified as in (1) and in addition $[S(\lambda) : L(\lambda)] = 1$ for all $\lambda \in \Lambda$, we say that $(A, \preceq)$ is *quasi-hereditary*.

Observe that if $(A, \preceq)$ is standardly stratified, $(A, \preceq')$ is also standardly stratified, for any extension $\preceq'$ of $\preceq$.

2.7.5. In case the order is total, Definitions 2.7.1 and 2.7.4 agree. For (5), this is [CPS1, Theorem 3.6] and for (1), this is [CPS2, Theorem 2.2.3]. For (3), the equivalence follows from the case (1) of standard stratifications, as going from (1) to (3) in both Definitions 2.7.1 and 2.7.4 only corresponds to restricting from partial quasi-orders to partial orders, see Remark 2.7.3. The equivalence of the definitions for (4) follows from the combination of [KM, Corollary 3] and [Dl, Theorem 5]. The equivalence of the definitions for (2) follows from [Fr, Proposition 7]. Also for orders which are not total, one can define stratified algebras in terms of chains of ideals, by considering coarser chains.

2.7.6. The modules $S(\lambda)$ in Definition 2.7.4 have simple top $L(\lambda)$ and are known as *standard modules*. They satisfy $S(\lambda) \cong Ae_\lambda / J_{j-1}e_\lambda$ with j the largest such that $J_{j-1}L(\lambda) = 0$. The modules $\overline{S}(\lambda)$ are *proper standard modules* and are given by $S(\lambda) / (Ae_\lambda \text{rad} S(\lambda))$.

2.8. Faithful covers and standard systems. Rouquier introduced notions of *faithfulness* of *quasi-hereditary covers*, see [Ro, Definition 4.37]. We extend this to standardly stratified algebras. Consider a cover C as in Section 2.4(a), which is standardly stratified. Such a cover is called *j-faithful* if the functor $F = \text{Hom}_C(Ce, -)$ induces isomorphisms

$$(2.5) \quad \text{Ext}_{C_0}^i(M, N) \xrightarrow{\sim} \text{Ext}_{C_0}^i(FM, FN), \quad \forall 0 \leq i \leq j,$$

for all modules M, N admitting a filtration with sections given by proper standard modules. Important examples of quasi-hereditary 1-covers are the q-Schur algebras of the Hecke algebra, by [HN] and the analogue of [DJM, DuS] in type B , by [Ro, Theorem 6.6].

Consider an abelian category $\mathcal{C}$ and a partially ordered set $(S, \leq)$. As in [DR2, Section 3] or [HHKP, Definition 10.1], a *standard system in $\mathcal{C}$ for S* is a set of objects $\{\Theta(p) \mid p \in S\}$ in $\mathcal{C}$, such that for all $p, q \in S$:

- (1) $\text{End}_{\mathcal{C}}(\Theta(p))$ is a division ring;
- (2) $\text{Hom}_{\mathcal{C}}(\Theta(p), \Theta(q)) = 0$ unless $p \leq q$;
- (3) $\text{Ext}_{\mathcal{C}}^1(\Theta(p), \Theta(q)) = 0$ unless $p < q$.

For a quasi-hereditary algebra $(A, \leq)$, the standard modules $\{S(\lambda), \lambda \in \Lambda\}$ form a standard system in $A\text{-mod}$ for $(\Lambda, \leq)$, see e.g. [DR2, Lemmata 1.2, 1.3 and 1.6].

2.9. Standardly based algebras and Schur algebras. There is a close connection, and large overlap, between quasi-hereditary algebras and *cellular algebras*, see [GL, Remark 3.10], [KX1, Proposition 4.1 and Corollary 4.2], [KX2, Theorem 1.1] and [Ca, Theorem 1.1]. Moreover, an interesting concept of “cellularly stratified” algebras was recently introduced in [HHKP], which combines properties of cellular and stratified algebras. Comparing or combining properties of cellular and stratified algebras is often complicated by the involutive anti-automorphism ι in the definition of a cell datum in [GL, Definition 1.1]. Omitting ι in the definition leads to the concept of standardly based algebras.

We use a reformulation of [DuR, Definition 1.1.2]. A *standardly based structure* of an algebra A is a poset L , with two-sided ideals $A^{\geq p}$ for each $p \in L$, such that for all $p, q \in L$

- $A^{\geq p} \supseteq A^{\geq q}$ if $p < q$;
- $A^{\geq p}/A^{>p} \cong W(p) \otimes_{\mathbb{k}} W'(p)$, with $A^{>p} := \cup_{q>p} A^{\geq q}$, as A -bimodules for a left module $W(p)$ and right module $W'(q)$.

The modules $W(p)$ will be referred to as the *cell modules* of A .

Remark 2.9.1. A standardly based structure of an algebra is preserved under Morita equivalences, see [YL, Section 3], meaning that the ideals are naturally linked. In particular, the cell modules are mapped to the corresponding cell modules.

By [DuR, Theorem 2.4.1], Λ can be naturally identified with a subset of L . We consider Λ then as a poset for the inherited partial order from L . For $\lambda \in \Lambda \subset L$, we have furthermore that $\text{Top}W(\lambda) = L(\lambda)$ and

$$[W(\lambda) : L(\lambda)] = 1 \quad \text{and} \quad [W(\lambda) : L(\mu)] = 0 \quad \text{unless } \mu \leq \lambda.$$

By [DuR, Proposition 2.4.4], every indecomposable projective module $P(\lambda)$ with $\lambda \in \Lambda$ has a filtration with sections given by modules $W(p)$ such that

$$(P(\lambda) : W(\lambda)) = 1 \quad \text{and} \quad (P(\lambda) : W(p)) = 0 \quad \text{unless } p \geq \lambda.$$

Hence if $L = \Lambda$, the standardly based algebra is quasi-hereditary with standard modules $W(\lambda)$. Conversely, over an algebraically closed field, every quasi-hereditary algebra is standardly based for $L = \Lambda$ (as posets) and $W(\lambda) = S(\lambda)$, by [DuR, Theorem 4.2.3].

2.9.2. *Example.* Consider $A := \mathbb{k}[x]/(x^t - 1)$ for a field $\mathbb{k}$ such that $x^t - 1$ splits as

$$x^t - 1 = \prod_{i=1}^t (x - \omega_i),$$

for not necessarily distinct $\omega_i \in \mathbb{k}$. Then consider $L := \{1, 2, \dots, t\}$ with usual order and

$$a_k = \prod_{i=1}^{k-1} (x - \omega_i) \quad \text{for } 2 \leq k \leq t \quad \text{and } a_1 = 1.$$

The ideals $A^{\geq i} = \text{Span}\{a_i, a_{i+1}, \dots, a_t\}$ give a standardly based structure of A . A is even cellular for involution $\iota = \text{id}_A$ the identity, by [DuR, Lemma 1.2.4], or [GL, Example 1.3].

2.9.3. *Example.* For any field $\mathbb{k}$, the group algebra $\mathbb{k}\mathbb{S}_t$ of the symmetric group $\mathbb{S}_t$ on t symbols is cellular and hence standardly based for $L = \{\lambda \vdash t\}$, equipped with the partial order obtained by reversing the dominance order [Ja, 1.4.6] on partitions, see [GL, Example 1.2]. The ideals are obtained from the Kazhdan-Lusztig basis (expressed in terms of the Robinson-Schensted correspondence) of the Hecke algebra specialised at $q = 1$ and the cell modules are the Specht modules of [Ja, Section 7.1]. The algebra $\mathbb{k}\mathbb{S}_t$ is then cellular for the involution ι , which is the linearisation of the inversion on the group $\mathbb{S}_t$.

The following definition is essentially [HHKP, Definition 12.1], see also [HK, Ro].

Definition 2.9.4. A *Schur algebra* of a standardly based algebra A is a quasi-hereditary 1-cover $(C, \leq)$, such that $(\Lambda_C, \leq) = (L_A, \leq)$ and $F(S(p)) \cong W(p)$ for all $p \in L_A$.

By [Ro, Corollary 4.46], a Schur algebra of a standardly based algebra is unique, up to Morita equivalence, if it exists.

Part I. General theory

3. PRE-BORELIC PAIRS

In this section we introduce the notion of pre-Borelic pairs for arbitrary algebras. These will lead to Borelic pairs in the next section, where we will also prove that these contain König's notion of exact Borel subalgebras for quasi-hereditary algebras as a special case.

3.1. **Definition and properties.** Fix an algebra A .

Definition 3.1.1. A pair (B, H) of subalgebras $H \subset B \subset A$ forms a *pre-Borelic pair* of A if there exists a two-sided ideal B_+ in B , with $B = H \oplus B_+$, such that

- (I) A is projective as a right B -module and B is projective as a left H -module;
- (II) taking B_+ -invariants yields an equivalence of categories between simple A -modules and simple H -modules;
- (III) the ideal B_+ is contained in the Jacobson radical $\text{rad} B$ of B .

3.1.2. Label the set of isomorphism classes of simple A -modules by Λ as in Section 2.3. By (II), we can use the same set for H . We use the notation $L(\lambda)$ (resp. $L^0(\lambda)$) for the corresponding simple modules over A (resp. H), and hence

$$(3.1) \quad L^0(\lambda) \cong L(\lambda)^{B_+}.$$

By assumption (III), we also have a one-to-one correspondence between simple B -modules and simple H -modules, so

$$(3.2) \quad \Lambda = \Lambda_A = \Lambda_B = \Lambda_H.$$

We use the same notation $L^0(\lambda)$, for the simple B -module with trivial B_+ -action defined as the inflation of the H -module $L^0(\lambda)$. For each e_λ , there is an idempotent $e_\lambda^0 \in H \subset B \subset A$, primitive in H (but generally not in A) such that $e_\lambda^0 L^0(\lambda) \neq 0$.

3.1.3. For any H -module N , interpreted as a B -module with trivial B_+ -action, we define

$$\Delta_N := \text{Ind}_B^A N = A \otimes_B N.$$

Its fundamental property is

$$(3.3) \quad \text{Hom}_A(\Delta_N, M) \cong \text{Hom}_H(N, M^{B_+}),$$

for all $M \in A\text{-mod}$, as follows from Frobenius reciprocity.

Lemma 3.1.4. *Consider a pre-Borelic pair (B, H) of A . For any $N \in H\text{-mod}$ with simple top $L^0(\lambda)$ for some $\lambda \in \Lambda$, we have*

$$\text{Top} \Delta_N \cong L(\lambda).$$

Proof. Equations (3.3) and (3.1) imply that, for $\lambda, \lambda' \in \Lambda$, we have

$$\dim \text{Hom}_A(\Delta_N, L(\lambda')) = \dim \text{Hom}_H(N, L^0(\lambda')) = \delta_{\lambda, \lambda'},$$

proving the claim. $\square$

Lemma 3.1.5. *Given a pre-Borelic pair (B, H) , define $P_\lambda = Ae_\lambda^0$, for any $\lambda \in \Lambda$. Then*

$$P_\lambda \cong \bigoplus_{\mu \in \Lambda} P(\mu)^{\oplus c_\mu^\lambda} \quad \text{with} \quad c_\mu^\lambda := [\text{Res}_B^A L(\mu) : L^0(\lambda)].$$

Proof. This follows from $A \otimes_B Be_\lambda^0 \cong Ae_\lambda^0$ and Frobenius reciprocity. $\square$

3.1.6. We introduce the modules

$$(3.4) \quad \Delta(\lambda) = \Delta_{He_\lambda^0} = A \otimes_B He_\lambda^0.$$

Consider an algebra A with pre-Borelic pair (B, H) . We say that an A -module N has a Δ -flag $\mathcal{M}$ if there is a $d \in \mathbb{N}$ and submodules M_i , $0 \leq i \leq d$:

$$(3.5) \quad N = M_0 \supset M_1 \supset M_2 \supset \cdots \supset M_{d-1} \supset M_d = 0,$$

such that M_i/M_{i+1} is in $\text{add}(\Delta)$ with $\Delta = \bigoplus_{\lambda \in \Lambda} \Delta(\lambda)$. For such a Δ -flag $\mathcal{M}$, we introduce

$$(N : \Delta(\lambda))_{\mathcal{M}} := \sum_{i=0}^{d-1} \dim \text{Hom}_A(M_i/M_{i+1}, L(\lambda)).$$

By definition and Lemma 3.1.4, we then have

$$(3.6) \quad [N] = \bigoplus_{\mu \in \Lambda} (N : \Delta(\mu))_{\mathcal{M}} [\Delta(\mu)],$$

in the Grothendieck group $G_0(A)$ of A -mod.

3.2. A special case: idempotent graded algebras. For an algebra A , we choose a decomposition

$$(3.7) \quad 1_A = e_0^* + e_1^* + \cdots + e_n^*,$$

where e_i^* are mutually orthogonal, but not necessarily primitive, idempotents. For notational convenience, we allow some of these idempotents to be zero. We consider a $\mathbb{Z}$ -grading on A :

$$(3.8) \quad A_j = \bigoplus_i e_i^* A e_{i+j}^*.$$

We call this the *idempotent grading* associated to the (ordered) choice of e_i^* in (3.7) and set

$$A_+ := \bigoplus_{j>0} A_j \quad \text{and} \quad A_- := \bigoplus_{j<0} A_j.$$

Definition 3.2.1. A $\mathbb{Z}_+$ -graded subalgebra of an idempotent graded algebra A is a graded subalgebra B such that $B_j = 0$ for $j < 0$. For such B set

$$H := B_0 = \bigoplus_{i \in \mathbb{N}} e_i^* B e_i^* \quad \text{and} \quad B_+ := \bigoplus_{j>0} B_j.$$

A $\mathbb{Z}_+$ -graded subalgebra B is *complete* if

$$(3.9) \quad A = B \oplus A_- B.$$

Note that any subalgebra B satisfying (3.9) is automatically $\mathbb{Z}_+$ -graded. Furthermore, it is easy to check that equation (3.9) is equivalent to

$$(3.10) \quad A = B \oplus A_- A.$$

We will use freely the fact that, as follows from the definitions, any module M , over A , B or H , is automatically a $\mathbb{Z}$ -graded module, by setting

$$M_{-j} := e_j^* M, \quad \text{for} \quad j \in \{0, 1, \dots, n\}.$$

Lemma 3.2.2. *Consider a complete $\mathbb{Z}_+$ -graded subalgebra B of an idempotent graded algebra A . If A is projective as a right B -module and B projective as a left H -module, then (B, H) is a pre-Borelic pair.*

Proof. Condition (I) in Definition 3.1.1 is given.

Condition (II) is the requirement that $\{L(\lambda)^{B_+}, \lambda \in \Lambda\}$ be a complete set of non-isomorphic simple H -modules. Simplicity of the $\mathbb{Z}$ -graded A -module $L(\lambda)$ implies that $L(\lambda)^{A_+}$ is the highest degree component and a simple A_0 -module. Equation (3.9) implies that $A_0 = H \oplus (A_- A_+)_0$, so the restriction of $L(\lambda)^{A_+}$ to an H -module remains simple. Equation (3.9) implies $A_+ \subset A B_+$, from which it follows that $L(\lambda)^{B_+} = L(\lambda)^{A_+}$ as H -modules. Hence,

$$(3.11) \quad L^0(\lambda) := L(\lambda)^{B_+}$$

is a simple H -module.

Now we consider an arbitrary simple H -module L^0 , which we can interpret as a simple B -module contained in one degree, say $-i$ for $i \in \mathbb{N}$, so $L^0(\lambda) = e_i^* L^0(\lambda)$. The A -module $M := \text{Ind}_B^A L^0$ satisfies $e_j^* M = 0$ unless $j \geq i$, with $e_i^* M \cong L^0(\lambda)$, by equation (3.9). It thus follows that $M \neq 0$, and that any proper submodule S of M satisfies $e_i^* S = 0$. Hence, taking the union of all proper submodules yields the unique maximal submodule. We conclude that

$\text{Ind}_B^A L^0$ has simple top. It follows from Frobenius reciprocity that this top is $L(\lambda)$ if and only if $L^0 \cong L^0(\lambda)$. This implies that the simple modules in (3.11) exhaust all simple H -modules and that $L^0(\lambda) \cong L^0(\mu)$ implies $L(\lambda) \cong L(\mu)$, which proves condition (II).

Condition (III) is immediate from the grading. This concludes the proof. $\square$

The lemma motivates the following definition.

Definition 3.2.3. Let A be an idempotent graded algebra with complete $\mathbb{Z}_+$ -graded subalgebra B . If A_B and ${}_H B$ are projective with $H := B_0$, we call B a *graded pre-Borelic algebra*.

The reason we just work with the algebra B for graded pre-Borelic subalgebras, instead of a pair, is that $H := B_0$ is defined through the grading on B .

Remark 3.2.4. An idempotent grading on an algebra A implies a chain of idempotent ideals. Set $f_k := \sum_{j < k} e_j^*$, for $1 \leq k \leq n+1$ and $J_k := Af_k A$, leading to the chain

$$(3.12) \quad 0 = J_0 \subset J_1 \subset \cdots \subset J_n \subset J_{n+1} = A.$$

Using the chain (3.12), we can rewrite equation (3.10) as

$$(3.13) \quad e_j^* A = e_j^* B \oplus e_j^* J_j, \quad \text{for all } 0 \leq j \leq n.$$

In particular, $e_j^* A e_j^* = e_j^* H \oplus e_j^* J_j e_j^*$, and for the algebras in Remark 2.6.2 we find

$$(3.14) \quad A^{(i)} = f_{i+1}(A/J_i)f_{i+1} = e_i^*(A/J_i)e_i^* \cong H e_i^*.$$

The following theorem shows that a graded pre-Borelic algebra implies an exact standard stratification. This will also be proved by a more general approach in Section 4.4.

Theorem 3.2.5. *Consider an algebra A with idempotent grading (3.8).*

- (1) *If A admits a graded pre-Borelic algebra B , then (3.12) is an exactly standardly stratifying chain and if H is quasi-local (resp. semisimple), the chain is properly stratifying (resp. quasi-heredity).*
- (2) *Assume that the chain (3.12) is exactly standardly stratifying. Then a subalgebra $B \subset A$ is a graded pre-Borelic subalgebra if and only if B satisfies equation (3.13), i.e. B is a complete $\mathbb{Z}_+$ -graded subalgebra.*

Remark 3.2.6. The theorem indicates that our methods might be well-suited to extend to the setting of [Kl]. With notation introduced in Appendix A.2, Theorem 3.2.5(1) can be paraphrased as: If an idempotent graded algebra A has graded pre-Borelic subalgebra B with $H \in \mathcal{B}$, then A is $\mathcal{B}$ -properly stratified, for $\mathcal{B}$ one of the classes of algebras $\mathcal{D}$, $\mathcal{L}$ or $\mathcal{S}$.

We start the proof of Theorem 3.2.5 with the following lemma.

Lemma 3.2.7. *Let A be an idempotent graded algebra with complete $\mathbb{Z}_+$ -graded subalgebra B . Then ${}_H B$ is projective if and only if the left $A^{(j)}$ -module $f_{j+1}A/J_j$ is projective for $0 \leq j \leq n$.*

Proof. By (3.14), we have $H e_j^* \cong A^{(j)}$ and by (3.13), $e_j^* B \cong f_{j+1}(A/J_j)$, which implies the statement. $\square$

Proof of Theorem 3.2.5. First we prove part (1). As A_B is projective, we have

$$A_B \cong \bigoplus_{j, \tau} e_{j, \tau} B,$$

for some idempotents $e_{j,\tau}$ of B satisfying $e_k^* e_{j,\tau} = \delta_{jk} e_{j,\tau}$. As B is $\mathbb{Z}_+$ -graded we find $Ae_0^* \cong \oplus_{\tau} e_{0,\tau} H$ and consequently $Ae_0^* B \cong \oplus_{\tau} e_{0,\tau} B$. By equation (3.13), we have $e_0^* B = e_0^* A$, so

$$(Ae_0^* A)_A \cong \bigoplus_{\tau} e_{0,\tau} A,$$

which implies that J_1 is projective as a right A -module. Furthermore, the above shows that J_1 is a direct summand of A as a right B -module. Hence A/J_1 is projective as a right B -module and we can proceed iteratively to prove that J_{j+1}/J_j is projective as a right A/J_j -module. Lemma 3.2.7 implies that $f_{j+1}A/J_j$ is a projective left $A^{(j)}$ -module. It then follows from Lemma A.1.2 that J_{j+1}/J_j is also projective as a left A/J_j -module.

Now we prove part (2). A graded pre-Borelic subalgebra B is complete $\mathbb{Z}_+$ -graded by definition, so we prove the other direction of the claim. Consider a complete $\mathbb{Z}_+$ -graded subalgebra B . Lemmata 3.2.7 and A.1.2(1) then imply that ${}_H B$ is projective.

Now we claim that for an exact standard stratification, the right A -module J_i is contained in $\text{add}(f_i A)$. The right A -module J_i is the union of the images of all morphisms from modules in $\text{add}(f_i A)$ to A_A . The claim then follows from the fact that an exact (standard) stratification of A implies a standard stratification of A^{op} , so we can use the arguments in the proof of [CPS2, Theorem 2.2.3].

Finally, we use the above claim to prove that A_B is projective. By equation (3.13) we have $e_0^* A = e_0^* B$. We use this as the first step to prove by induction on i that $e_i^* A$ is a projective right B -module. Indeed, by equation (3.13) and the above claim, the right B -module $e_i^* A$ decomposes as $e_i^* B$ and the direct sum of A -direct summands (and hence B -direct summands) of $e_j^* A$ for $j < i$. $\square$

The following observation is an immediate consequence of equation (3.13).

Lemma 3.2.8. *Consider an idempotent graded algebra A with graded pre-Borelic algebra B , and an $He_i^* \cong A^{(i)}$ -module M . We have*

$$A \otimes_B M \cong (A/J_i) f_{i+1} \otimes_{A^{(i)}} M.$$

4. BORELIC PAIRS

4.1. Definition. Consider an algebra A with pre-Borelic pair (B, H) .

4.1.1. We define an equivalence relation $\sim_H$ on Λ , where $\lambda \sim_H \mu$ if and only if $L^0(\lambda)$ and $L^0(\mu)$ are in the same indecomposable block of $H\text{-mod}$. This equivalence relation is hence trivial if and only if $e_\lambda^0 H e_\mu^0 = 0$ for all $\mu \neq \lambda$. In particular, the equivalence relation is trivial when $B_+ = \text{rad} B$. A partial quasi-order $\preceq$ on Λ is said to be *H -compatible* if

$$\lambda \sim_H \mu \Rightarrow \lambda \preceq \mu,$$

where we recall that $\lambda \sim \mu$ stands for $\lambda \preceq \mu$ and $\mu \preceq \lambda$.

Definition 4.1.2. Consider an H -compatible partial quasi-order $\preceq$ on Λ . We say that (B, H) is a *Borelic pair* of $(A, \preceq)$ if, for all $\lambda, \mu \in \Lambda$, we have

- (1) $[B_+ e_\lambda^0 : L^0(\mu)] = 0$ unless $\mu \succ \lambda$;
- (2) $[\Delta(\lambda) : L(\mu)] = 0$ unless $\mu \preceq \lambda$;
- (3) $[\text{Res}_B^A L(\mu) : L^0(\lambda)] = 0$ unless $\mu = \lambda$ or $\lambda \prec \mu$.

A Borelic pair (B, H) is called *exact* if $[A \otimes_B L^0(\lambda) : L(\lambda)] = 1$, for all $\lambda \in \Lambda$.

4.2. Properties. Now we start exploring the properties of Borelic pairs. Whenever a partial quasi-order $\preccurlyeq$ on Λ is considered in relation to a pre-Borelic pair (B, H) of A , we assume it to be H -compatible.

Lemma 4.2.1. *Consider an algebra $(A, \preccurlyeq)$ with Borelic pair (B, H) , then*

$$\mathrm{Hom}_A(\Delta(\lambda), \Delta(\mu)) = 0 \quad \text{unless } \lambda \preccurlyeq \mu.$$

Proof. This is a consequence of Definition 4.1.2(2) and Lemma 3.1.4. $\square$

Lemma 4.2.2. *Consider an algebra $(A, \preccurlyeq)$ with pre-Borelic pair (B, H) satisfying Definition 4.1.2(1), then the module $P_\lambda = A \otimes_B Be_\lambda^0$ admits Δ -flags. In particular, the module K defined through the short exact sequence*

$$(4.1) \quad 0 \rightarrow K \rightarrow P_\lambda \rightarrow \Delta(\lambda) \rightarrow 0,$$

has a Δ -flag $\mathcal{M}$ with

$$(K : \Delta(\mu))_{\mathcal{M}} = 0 \quad \text{unless } \mu \succ \lambda.$$

Proof. The short exact sequence (4.1) is induced from the short exact sequence

$$0 \rightarrow B_+e_\lambda^0 \rightarrow Be_\lambda^0 \rightarrow He_\lambda^0 \rightarrow 0,$$

of B -modules, through the exact functor $A \otimes_B -$, see Definition 3.1.1(I).

Now we claim that the B -module $B_+e_\lambda^0$ has a filtration with sections given by the B -modules He_μ^0 with $\mu \succ \lambda$. Take $\mu \in \Lambda$ such that there is no $\mu' \in \Lambda$ with $\mu' \succ \mu$. Then the union M of all images of morphisms $Be_\nu^0 \rightarrow B_+e_\lambda^0$ with $\nu \sim \mu$ forms a B -submodule with trivial B_+ action, by Definition 4.1.2(1). This B -submodule M is a direct summand as an H -module, since $\sim$ is H -compatible. Hence both the B -modules M and $B_+e_\lambda^0/M$ are projective as H -modules by Definition 3.1.1(I). Consequently M is a direct sum of B -modules He_ν^0 with $\nu \sim \mu$, and is only non-zero if $\nu \succ \lambda$ by Definition 4.1.2(1). One can continue this procedure by taking a maximal element in the ordered set Λ obtained by removing the equivalence class containing μ .

The result therefore follows from the fact that $\mathrm{Ind}_B^A -$ is exact by Definition 3.1.1(I). $\square$

Lemma 4.2.3. *Consider an algebra $(A, \preccurlyeq)$ with Borelic pair (B, H) . The modules $\Delta(\lambda)$ induce a $\mathbb{k}$ -basis $\{[\Delta(\lambda)] \mid \lambda \in \Lambda\}$ of $G_0(A)$.*

Proof. The indecomposable projective A -modules induce a basis $\{[P(\lambda)] \mid \lambda \in \Lambda\}$. The combination of Lemma 3.1.5 and Definition 4.1.2(3) imply that another basis is given by $\{[P_\lambda] \mid \lambda \in \Lambda\}$. Lemma 4.2.2 and equation (3.6) then imply that $\{[\Delta(\lambda)] \mid \lambda \in \Lambda\}$ also forms a basis. $\square$

By equation (3.6) and Lemma 4.2.3, the multiplicities $(N : \Delta(\mu))_{\mathcal{M}}$ coincide for all possible Δ -flags $\mathcal{M}$, if (B, H) is a pre-Borelic pair satisfying Definition 4.1.2(1). Hence, under that assumption, we leave out the reference to $\mathcal{M}$ in the multiplicities.

Lemma 4.2.4. *Consider an algebra $(A, \preccurlyeq)$ with Borelic pair (B, H) . We have*

$$\mathrm{Ext}_A^1(\Delta(\lambda), \Delta(\mu)) = 0 \quad \text{unless } \lambda \prec \mu.$$

Proof. Consider K as in Lemma 4.2.2. Elementary homological algebra implies a surjection

$$\mathrm{Hom}_A(K, \Delta(\mu)) \twoheadrightarrow \mathrm{Ext}_A^1(\Delta(\lambda), \Delta(\mu)).$$

By Lemmata 4.2.1 and 4.2.2, the left-hand side (and therefore the right-hand side) is zero unless there exists $\nu \in \Lambda$ with $\nu \succ \lambda$ and $\nu \preccurlyeq \mu$. This implies in particular that $\mu \succ \lambda$. $\square$

Lemma 4.2.5. *Consider an algebra $(A, \preceq)$ with Borelic pair (B, H) . If for an A -module M with Δ -flag, we have $[M : L(\nu)] = 0$ for all $\nu \in \Lambda$ with $\nu \succ \lambda$ and $[M : L(\lambda)] \neq 0$, then there exists an injection*

$$\Delta(\lambda') \hookrightarrow M,$$

for some $\lambda' \sim \lambda$. Moreover, for any such injection, the cokernel also has a Δ -flag.

Proof. We consider a filtration of M as in (3.5), where each section is indecomposable, so isomorphic to some $\Delta(\mu)$. The fact that $[\Delta(\mu) : L(\mu)] \neq 0$, by Lemma 3.1.4 and $[\Delta(\mu) : L(\nu)] = 0$ unless $\nu \preceq \mu$ by Definition 4.1.2(2), imply that there must be some i such that $M_i/M_{i+1} \cong \Delta(\lambda')$ for some $\lambda' \sim \lambda$ and that all other sections are of the form $\Delta(\mu)$ with $\mu \preceq \lambda \sim \lambda'$. Now we have a short exact sequence

$$0 \rightarrow M_{i+1} \rightarrow M_i \rightarrow \Delta(\lambda') \rightarrow 0,$$

where M_{i+1} thus has a filtration with sections given by $\Delta(\mu)$ with $\mu \preceq \lambda'$. Lemma 4.2.4 implies that this extension splits, which implies $\Delta(\lambda') \hookrightarrow M$.

For any μ , the number of times $\Delta(\mu)$ appears as a section of a fixed Δ -flag of M is determined by $[M]$ through equation (3.6). By the above reasoning, there is a module $\oplus_i \Delta(\lambda_i)$, with $\lambda_i \sim \lambda$, which injects into M , such that the cokernel has a Δ -flag, where there are no more sections of the form $\Delta(\lambda')$ for $\lambda' \sim \lambda$ appearing. The image of any such $\Delta(\lambda') \hookrightarrow M$ must be in that constructed submodule of M , concluding the proof. $\square$

Finally we show that one can replace Definition 4.1.2(3) by a different condition.

Lemma 4.2.6. *Consider an algebra $(A, \preceq)$ with pre-Borelic pair (B, H) satisfying conditions (1) and (2) in Definition 4.1.2. Then (B, H) is a Borelic pair if and only if for any $M \oplus M' \in A\text{-mod}$ with a Δ -flag, M and M' also have Δ -flags.*

Proof. Assume that (B, H) is a Borelic pair. Take some λ for which we can apply Lemma 4.2.5 to $M \oplus M'$. We have some $\Delta(\lambda') \hookrightarrow M \oplus M'$. If the image is not contained in either M or M' , the space of such morphisms must have higher dimension and by Lemma 4.2.5 each such morphism leads to a factor module with Δ -flag. Choosing an injection such that the image is contained in one of the two summands we hence obtain a direct sum of two modules with Δ -flag, where the length of the flag is one lower. This procedure can be repeated iteratively to prove that both M and M' have Δ -flags.

Now consider a pre-Borelic pair (B, H) , satisfying conditions (1) and (2) in Definition 4.1.2. Assume that for any $M \oplus M' \in A\text{-mod}$ with a Δ -flag, M and M' also have a Δ -flag. Lemmata 3.1.5 and 4.2.2 imply that every $P(\mu)$ has a Δ -flag. We claim that the same is true for the kernel K_0 of $P(\mu) \twoheadrightarrow \Delta(\mu)$. Consider K as in (4.1), with Δ -flag. It follows easily that K contains K_0 as a direct summand, hence K_0 has a Δ -flag with

$$(4.2) \quad (P(\mu) : \Delta(\mu)) \neq 0 \quad \text{for any } \mu \in \Lambda.$$

Using this observation we now prove condition (3) in Definition 4.1.2. Consider $\lambda \neq \mu$, such that $[\text{Res}_B^A L(\mu) : L^0(\lambda)] \neq 0$. Lemma 3.1.5 implies that $P(\mu)$ is a direct summand of P_λ . By equation (4.2), we then find that $(P_\lambda : \Delta(\mu)) \neq 0$. Lemma 4.2.2 thus implies that $\mu \succ \lambda$. Hence we find that also condition (3) is satisfied, so that (B, H) is Borelic. $\square$

Corollary 4.2.7. *Consider an algebra $(A, \preceq)$ with Borelic pair (B, H) . Then the kernel K_0 of $P(\lambda) \twoheadrightarrow \Delta(\lambda)$ has a Δ -flag with*

$$(K_0 : \Delta(\nu)) = 0 \quad \text{unless } \nu \succ \lambda.$$

4.3. Exact Borel subalgebras for exactly standardly stratified algebras. Before we present the properties of algebras with Borelic pairs we need to review the notion of exact Borel subalgebras.

An *exact Borel subalgebra of a quasi-hereditary algebra* $(A, \leq)$, as introduced by König in [Kö], is a subalgebra B satisfying the following three conditions from [KKO, Definition 2.2].

- (1) The simple modules of B are also labelled by Λ_A , so $\Lambda_A = \Lambda_B = \Lambda$ and we denote the simple B -modules by $\{L^0(\lambda) \mid \lambda \in \Lambda\}$. The only simple subquotients in the radical of the projective cover of $L^0(\lambda)$ are $L^0(\mu)$ with $\mu \succ \lambda$.
- (2) The functor $A \otimes_B -$ is exact.
- (3) There is an isomorphism $A \otimes_B L^0(\lambda) \cong S(\lambda)$ for each $\lambda \in \Lambda$.

For an algebra B with simple modules $L^0(\lambda)$ for $\lambda \in \Lambda$, choose a decomposition $1 = \sum_{\lambda \in \Lambda} \bar{e}_\lambda$, where $\{\bar{e}_\lambda\}$ is a set of mutually orthogonal idempotents, satisfying $\bar{e}_\mu L^0(\lambda) = 0$ unless $\mu = \lambda$.

Lemma 4.3.1. *An exact Borel subalgebra B of a quasi-hereditary algebra $(A, \leq)$ is Wedderburn. We have*

$$B = H \oplus \text{rad}B \quad \text{with } H := \bigoplus_{\lambda \in \Lambda} \bar{e}_\lambda B \bar{e}_\lambda.$$

Proof. As in [KM, Lemma 3] one proves that $\bar{e}_\lambda B \bar{e}_\lambda$ is a division algebra and hence H is semisimple. Furthermore $\bar{e}_\mu B \bar{e}_\lambda = 0$ unless $\mu \geq \lambda$, showing that

$$\bigoplus_{\mu \neq \lambda} \bar{e}_\lambda B \bar{e}_\mu = \bigoplus_{\mu \succ \lambda} \bar{e}_\lambda B \bar{e}_\mu \subset \text{rad}B.$$

The statement thus follows. $\square$

In [KM, Definition 1], the notion of exact Borel subalgebras was generalised to properly stratified algebras and it can be further generalised to exactly standardly stratified algebras.

Definition 4.3.2. An *exact Borel subalgebra of an exactly standardly stratified algebra* $(A, \preceq)$ is a subalgebra B such that the following four conditions are satisfied.

- (1) The simple modules of B are also labelled by Λ_A , so $\Lambda_A = \Lambda_B = \Lambda$ and we denote the simple B -modules by $\{L^0(\lambda) \mid \lambda \in \Lambda\}$. We have $\bar{e}_\mu B \bar{e}_\lambda = 0$ unless $\lambda \preceq \mu$.
- (2) The functor $A \otimes_B -$ is exact.
- (3) The algebra

$$H := \bigoplus_{\lambda \sim \mu} \bar{e}_\lambda B \bar{e}_\mu \cong B / \left(\bigoplus_{\mu \prec \lambda} \bar{e}_\lambda B \bar{e}_\mu \right)$$

is such that the left H -module ${}_H B$ is projective. The interpretation of H as a quotient algebra of B allows to interpret H -modules as B -modules by inflation.

- (4) There are isomorphisms $A \otimes_B L^0(\lambda) \cong \bar{S}(\lambda)$ and $A \otimes_B P^0(\lambda) \cong S(\lambda)$, with $P^0(\lambda)$ the H -projective cover of $L^0(\lambda)$, for each $\lambda \in \Lambda$.

Remark 4.3.3. There is no conflict between Definition 4.3.2 and the special case of quasi-hereditary algebras.

An exact Borel subalgebra B of a quasi-hereditary algebra $(A, \preceq)$ is an exact Borel subalgebra of $(A, \preceq)$, interpreted as an exactly standardly stratified algebra. This follows from Lemma 4.3.1.

Now consider an exact Borel subalgebra B of an exactly standardly stratified algebra $(A, \preceq)$ which happens to be quasi-hereditary. As $S(\lambda) \cong \bar{S}(\lambda)$, Definition 4.3.2(4) implies

that H is semisimple. It then follows that the multiplicity of $L^0(\lambda)$ in its projective cover over B is 1. Hence B is an exact Borel subalgebra of $(A, \preceq)$ as a quasi-hereditary algebra.

Remark 4.3.4. If $(A, \preceq)$ is properly stratified, Definition 4.3.2 is almost identical to [KM, Definition 1]. The latter also requires projectivity of the *right* H -module B . This is equivalent to demanding that B itself is also properly stratified, see [KM, Proposition 2].

4.4. Algebras with Borelic pairs.

Theorem 4.4.1. *If an algebra A admits a Borelic pair (B, H) for an H -compatible partial quasi-order $\preceq$, then $(A, \preceq)$ is standardly stratified. If moreover $\preceq$ is a partial order (implying that H is quasi-local) then $(A, \preceq)$ is strongly standardly stratified.*

Theorem 4.4.2. *Consider an algebra $(A, \preceq)$ for some partial quasi-order $\preceq$.*

- (1) *Assume that $\preceq$ is a partial order. The following statements are equivalent, for any subalgebra B of A :*
 - $(B, B/\text{rad}B)$ is an exact Borelic pair.
 - $(A, \preceq)$ is quasi-hereditary and B is an exact Borel subalgebra.
- (2) *Assume that $\preceq$ is a partial order. The following statements are equivalent, for any subalgebras $H \subset B$ of A :*
 - (B, H) is an exact Borelic pair.
 - $(A, \preceq)$ is properly stratified and B is an exact Borel subalgebra.
- (3) *The following statements are equivalent, for any subalgebras $H \subset B$ of A :*
 - (B, H) is an exact Borelic pair.
 - $(A, \preceq)$ is exactly standardly stratified and B is an exact Borel subalgebra.

Proof of Theorem 4.4.1. Consider $(A, \preceq)$ with Borelic pair (B, H) . That the modules $S(\lambda) := \Delta(\lambda)$ satisfy Definition 2.7.4(1) follows from Definition 4.1.2(2) and Corollary 4.2.7. The same reasoning for the case where $\preceq$ is a partial order implies that $(A, \preceq)$ is strongly standardly stratified as in Definition 2.7.4(2). $\square$

Lemma 4.4.3. *If A is exactly standardly stratified with an exact Borel subalgebra B with subalgebra H as in Definition 4.3.2, then (B, H) is a pre-Borelic pair.*

Proof. We check the conditions in Definition 3.1.1. Condition 3.1.1(I) follows from 4.3.2(2) and (3). We define the two-sided ideal $B_+ := \oplus_{\lambda \not\sim \mu} \bar{e}_\lambda B \bar{e}_\mu = \oplus_{\lambda \succ \mu} \bar{e}_\lambda B \bar{e}_\mu$, for which we have $B = H \oplus B_+$. Condition 3.1.1(III) is clearly satisfied.

As the modules $S(\lambda)$ of 2.7.1(1) have simple top $L(\lambda)$, condition 4.3.2(4) leads, using Frobenius reciprocity, to

$$\delta_{\lambda, \mu} = \dim \text{Hom}_B(P^0(\lambda), L(\mu)) = \dim \text{Hom}_H(P^0(\lambda), L(\mu)^{B_+}) = [L(\mu)^{B_+} : L^0(\lambda)].$$

Condition 3.1.1(II) follows from this equation. $\square$

Proof of Theorem 4.4.2. It suffices to prove part (3). Part (2) is a special case and so is part (1) by remark 4.3.3.

Firstly, we assume that (B, H) is a Borelic pair. By Corollary 4.2.7 and Definition 4.1.2(2), it follows that A is standardly stratified as in Definition 2.7.4(1), with standard modules $S(\lambda) := \Delta(\lambda)$. Furthermore $S(\lambda)$ has a filtration where the sections are given by $\bar{S}(\mu) := A \otimes_B L^0(\mu)$ by Definition 3.1.1(I), with $\mu \sim_H \lambda$. By Definition 4.1.2 we have $[\bar{S}(\mu) : L(\mu)] = 1$, so $(A, \preceq)$ is exactly standardly stratified.

Now we prove that $(B, \preceq)$ is an exact Borel subalgebra, as in Definition 4.3.2. Condition 4.3.2(1) is satisfied by Definition 4.1.2(1) and condition 4.3.2(2) by 3.1.1(I). To prove the two remaining conditions in 4.3.2(3) and (4) we need to show that the subalgebra of B defined in 4.3.2(3), which we denote by H' , is the same as the subalgebra H in the pair (B, H) . Definition 4.1.2(1) implies that, as vector spaces, $H \subset H'$. The other inclusion follows from the fact that $\sim$ is assumed to be H -compatible.

Secondly, we assume that $(A, \preceq)$ is exactly standardly stratified and B is an exact Borel subalgebra. By Lemma 4.4.3, (B, H) is a pre-Borelic pair, so it remains to check the conditions in Definition 4.1.2. Condition 4.1.2(1) follows from 4.3.2(1) and 4.3.2(3). Condition 4.1.2(2) follows from Definitions 2.7.4(1) and 4.3.2(4). The extra condition for exact Borelic pairs follows from 4.3.2(4).

By the above, (B, H) is a pre-Borelic pair satisfying conditions (1) and (2) in Definition 4.1.2. From [Fr, Theorem 13], it follows in particular that a direct summand of a module with Δ -flag also has a Δ -flag. Condition 4.1.2(3) thus follows from Lemma 4.2.6. $\square$

4.5. A special case: graded Borelic subalgebras. For an idempotent graded algebra A as in Section 3.2, we have the corresponding chain (3.12) of idempotent ideals and hence an n -decomposition $\mathcal{Q}$ of $\Lambda = \sqcup_{i=0}^n \Lambda_i$, as in Section 2.6.3, with corresponding quasi-order and partial-order. We now introduce these directly.

Definition 4.5.1. The function $F_A : \Lambda \rightarrow \{0, 1, \dots, n\}$, where $F_A(\lambda)$ is the minimal $i \in \mathbb{N}$ such that there is an idempotent $f \in e_i^* A e_i^*$, equivalent to e_λ , then $\Lambda_i := \{\lambda \in \Lambda \mid F_A(\lambda) = i\}$.

- (1) The *idempotent quasi-order* is the total quasi-order $\preceq_{\mathcal{Q}}$ on Λ such that $\mu \preceq_{\mathcal{Q}} \lambda$ if and only if $F_A(\mu) \geq F_A(\lambda)$.
- (2) The *idempotent partial order* is the partial order $\leq_{\mathcal{Q}}$ on Λ such that $\mu <_{\mathcal{Q}} \lambda$ if and only if $F_A(\mu) > F_A(\lambda)$

Lemma 4.5.2. *For any $\lambda \in \Lambda$ we have $e_\lambda^0 \in H e_i^*$ with $i = F_A(\lambda)$. In particular, the partial quasi-order $\preceq_{\mathcal{Q}}$ is H -compatible.*

Proof. It follows from equation (3.11) that $L^0(\lambda)$ is a vector space contained in the highest degree of the graded vector space $L(\lambda)$. This implies that $e_i^* L^0(\lambda) \neq 0$ for the lowest i for which there is an idempotent e equivalent to e_λ such that $e e_i^* = e$.

Hence, for that minimal i , which is $F_A(\lambda)$, we have $e_\lambda^0 e_i^* = e_\lambda^0$. $\square$

For a graded pre-Borelic algebra, we also set

$$\overline{\Delta}(\lambda) := \Delta_{L^0(\lambda)} = A \otimes_B L^0(\lambda).$$

The notation is justified by the following proposition, which shows that a graded pre-Borelic algebra automatically implies an exact Borel pair.

Proposition 4.5.3. *Consider an algebra A with an idempotent grading. Let B be a graded pre-Borelic subalgebra with $H := B_0$.*

- (1) *The pair (B, H) is an exact Borelic pair for idempotent quasi-order $\preceq_{\mathcal{Q}}$.*
- (2) *If H is quasi-local, (B, H) is an exact Borelic pair for idempotent partial order $\leq_{\mathcal{Q}}$.*

Combining this with Theorem 4.4.2 yields the following corollary, which also gives an alternative proof of Theorem 3.2.5(1).

Corollary 4.5.4. *Consider an algebra A with an idempotent grading and graded pre-Borelic subalgebra B with $H := B_0$.*

- (1) Then $(A, \preceq_{\mathcal{Q}})$ is exactly standardly stratified with exact Borel subalgebra B .
- (2) If H is quasi-local, $(A, \leq_{\mathcal{Q}})$ is properly stratified with exact Borel subalgebra B .
- (3) If H is semisimple, $(A, \leq_{\mathcal{Q}})$ is quasi-hereditary with exact Borel subalgebra B .

Proof of Proposition 4.5.3. First observe that the graded A -module $\overline{\Delta}(\lambda)$ lives in degrees $\{-j, -j-1, \dots\}$ with $j = F_A(\lambda)$, by Lemma 4.5.2 and equation (3.9). Moreover, the degree $-j$ -component is precisely $L^0(\lambda)$. Each simple module $L(\mu)$ lives similarly in degree $\{-k, -k-1, \dots\}$ with $k = F_A(\mu)$. Therefore we find that

$$(4.3) \quad [\overline{\Delta}(\lambda) : L(\lambda)] = 1 \quad \text{and} \quad [\overline{\Delta}(\lambda) : L(\mu)] = 0 \text{ unless } \mu \leq_{\mathcal{Q}} \lambda.$$

Hence the corresponding condition in Definition 4.1.2 is satisfied.

Now we focus on the proof of part (1). By the above we only need to check the conditions in Definition 4.1.2. We have

$$[B_+ e_{\lambda}^0 : L^0(\mu)] = \sum_{j>0} \dim e_{\mu}^0 B_j e_{\lambda}^0.$$

By Lemma 4.5.2 we find that $e_{\mu}^0 B_j e_{\lambda}^0 \neq 0$ for $j > 0$ implies that $\mu \succ_{\mathcal{Q}} \lambda$, proving condition 4.1.2(1). The module $\Delta(\lambda) = A \otimes_B H e_{\lambda}^0$ has a filtration with sections $\overline{\Delta}(\mu) = A \otimes_B L^0(\mu)$ with $\mu \sim_H \lambda$, by Definition 3.1.1(I). Hence, by equation (4.3), $[\Delta(\lambda) : L(\mu)]$ implies there is a $\nu \in \Lambda$ with

$$\mu \leq_{\mathcal{Q}} \nu \sim_H \lambda \quad \text{and hence} \quad \mu \preceq_{\mathcal{Q}} \lambda.$$

So condition 4.1.2(2) for $\preceq_{\mathcal{Q}}$ is satisfied. Condition 4.1.2(3) follows similarly from the grading.

For the proof of part (2) we only need to observe $<_{\mathcal{Q}}$ and $\prec_{\mathcal{Q}}$ are identical and that, since H is quasi-local, now $\Delta(\lambda)$ has a filtration where each section is isomorphic to $\overline{\Delta}(\lambda)$. $\square$

Remark 4.5.5. By the proofs in Sections 3.2 and 4.5, we find that Corollary 4.5.4 remains true for a $\mathbb{Z}_+$ -graded subalgebra B , such that A_B and ${}_H B$ are projective, and for which

$$A_+ \subset AB_+ \quad \text{and} \quad A_0 = H \oplus (AB_+)_0.$$

However, in all examples of quasi-hereditary algebras with exact Borel subalgebras we will encounter in Part II, the stronger condition (3.9) is satisfied.

5. BASE STRATIFIED ALGEBRAS

5.1. Definition. The following definition is inspired by [HHKP, Propositions 5.1 and 5.2].

Definition 5.1.1. A *base-stratifying ideal* in an algebra A is an idempotent ideal $J = AeA$ which is exactly standardly stratifying, i.e. ${}_A J$ and J_A are projective, see 2.6.1, and such that eAe is standardly based, see Section 2.9.

5.1.2. Example. By Corollary 1, *any* exactly standardly stratifying ideal can be given the structure of a base-stratifying ideal, when working over an algebraically closed field.

Remark 5.1.3. By equation (2.3) and Remark 2.9.1, the definition above does not intrinsically depend on the choice of the idempotent e .

We use Definition 5.1.1 to introduce a generalisation of the concept of “cellularly stratified algebra” of [HHKP, Definition 2.1], see also [HHKP, Section 5].

Definition 5.1.4. Consider an algebra A with an n -decomposition $\mathcal{Q}$: $\Lambda = \sqcup_{i=0}^n \Lambda_i$. We consider the corresponding chain of idempotent ideals $\{J_i \mid 0 \leq i \leq n+1\}$ of 2.6.3. Then $(A, \mathcal{Q})$ is *base stratified* if each J_i/J_{i-1} is a base-stratifying ideal in A/J_{i-1} .

Remark 5.1.5. A trivial example of a base stratified algebra is any standardly based algebra with the 0-decomposition of Λ . It is hence essential to keep track of the decomposition $\mathcal{Q}$ of a base stratified algebra $(A, \mathcal{Q})$ in order to make non-trivial statements.

5.1.6. For a base stratified algebra $(A, \mathcal{Q})$, we make a choice of idempotents $\{f_l \mid 1 \leq l \leq n+1\}$ as in Remark 2.6.5 and consider the corresponding algebras $A^{(l)}$. We consider the set $L := \sqcup_{i=0}^n L_i = \{(i, p) \mid 1 \leq i \leq n, p \in L_i\}$, with L_i the poset of the standardly based algebra $A^{(i)}$. We make L into a poset by setting $(i, p) \leq (j, q)$ if $i > j$ or $i = j$ and $p \leq q$, so we take the lexicographic ordering up to reversal of the order on natural numbers.

Theorem 5.1.7. *A base stratified algebra $(A, \mathcal{Q})$ is standardly based for poset L .*

Proof. We use induction on n , where $\mathcal{Q}$ is an n -decomposition. If $n = 0$, there is nothing to prove. Assume the statement is true for $n - 1$ and consider the case n . Clearly the algebra $\tilde{A} := A/J_1$ is base stratified and hence standardly based for the poset $L' = \sqcup_{i=1}^n L_i$. We fix corresponding ideals $\tilde{A}^{\geq(l,p)}$ with $1 \leq l \leq n$ and $p \in L_l$.

We set $K := A^{(0)} = f_1 A f_1$. By Definition 5.1.1 and Lemma A.1.2 we have

$$J_1 \cong A f_1 \otimes_{f_1 A f_1} f_1 A \cong A f_1 \otimes_K K \otimes_K f_1 A.$$

By assumption, K is standardly based and we set

$$A^{\geq(0,p)} := A f_1 \otimes_K K^{\geq p} \otimes_K f_1 A,$$

for all $p \in L_0$. The fact that the quotients $A^{\geq(0,p)} / A^{>(0,p)}$ satisfy the desired properties then follows from the exactness of the functors $A f_1 \otimes_K -$ and $- \otimes_K f_1 A$ in Lemma A.1.2.

The set of ideals of A is then completed by taking $A^{(l,p)}$ to be an arbitrary ideal in A containing J_1 with $A^{(l,p)} / J_1 = \tilde{A}^{\geq(l,p)}$ for $l > 0$ and $p \in L_l$. $\square$

Corollary 5.1.8. *Maintain the notation and assumptions of Theorem 5.1.7. The cell modules of the standardly based algebra A satisfy*

$$W(i, p) \cong A f_{i+1} \otimes_{f_{i+1} A f_{i+1}} W_i^0(p),$$

with $W_i^0(p)$ the cell modules of $A^{(i)}$, regarded as $f_{i+1} A f_{i+1}$ -modules with trivial $f_{i+1} J_i f_{i+1}$ -action.

Remark 5.1.9. It follows from the considerations in Remark 2.6.2 that the cell module $W(i, p)$ of A does not depend on the actual choice of idempotents $\{f_i\}$.

Proposition 5.1.10. *Maintain the notation and assumptions of Theorem 5.1.7. For $0 \leq i \leq j \leq n$, $p \in L_i$, $q \in L_j$ and all $k \in \mathbb{N}$, we have*

$$\text{Ext}_A^k(W(i, p), W(j, q)) \cong \delta_{i,j} \text{Ext}_{A^{(i)}}^k(W_i^0(p), W_i^0(q)).$$

Proof. One immediately proves this by combining Corollary 5.1.8 with Lemma A.1.1. $\square$

This gives a very simple proof of the following analogue of [HHKP, Theorem 10.2].

Corollary 5.1.11. *Maintain the notation and assumptions of Theorem 5.1.7. The cell modules of A form a standard system for poset L if and only if, for each $0 \leq i \leq n$, the cell modules of $A^{(i)}$ form a standard system for poset L_i .*

Corollary 5.1.12. *A base stratified algebra $(A, \mathcal{Q})$ has standard modules $S(i, \lambda)$, properly standard modules $\overline{S}(i, \lambda)$ with $0 \leq i \leq n$ and $\lambda \in \Lambda_i$, and cell modules $W(i, p)$ with $0 \leq i \leq n$ and $p \in L_i$, such that for any fixed i ,*

- $S(i, \lambda)$ has a filtration where the sections are modules $W(i, p)$, $p \in L_i$ with $p \geq \lambda$;
- $W(i, p)$ has a filtration where the sections are modules $\overline{S}(i, \lambda)$, $\lambda \in \Lambda_i$.

Proof. The filtration of $S(i, \lambda)$ is induced from the filtration of the projective left $A^{(i)}$ -module using its cell modules. The filtration of $W(i, p)$ is induced from the Jordan-Hölder decomposition series of the $A^{(i)}$ -module $W_i^0(\lambda)$. $\square$

5.2. Base-Borelic pairs. Given an algebra A , we fix a decomposition of unity as in (3.7). Assume that we have a graded pre-Borelic algebra B as in Definition 3.2.3, with

$$H = \bigoplus_{i=0}^n e_i^* H e_i^* = \bigoplus_{i=0}^n H e_i^*.$$

This decomposition of H corresponds to an n -decomposition $\mathcal{Q}_H$ of $\Lambda = \Lambda_H = \Lambda_A$, where Λ_i consists of those $\lambda \in \Lambda$ satisfying $e_\lambda^0 e_i^* = e_\lambda^0$.

Theorem 5.2.1. *Consider an algebra A with a graded pre-Borelic algebra B such that $H = B_0$ is standardly based. Then $(A, \mathcal{Q}_H)$ is base stratified.*

Proof. This follows immediately from Theorem 3.2.5(1) and equation (3.14). $\square$

Corollary 5.2.2. *Consider an algebra A with a graded pre-Borelic algebra B such that H is standardly based. For each $0 \leq i \leq n$ consider the poset L_i (containing Λ_i) of the standardly based algebra He_i^* . Then A is standardly based for poset L defined in 5.1.6.*

Moreover, the cell modules over A form a standard system for L , if and only if the cell modules for He_i^* form a standard system for L_i , for each $0 \leq i \leq n$.

Proof. The first statement follows from the combination of Theorems 5.2.1 and 5.1.7. The claim about standard systems follows from Corollary 5.1.11 and Theorem 3.12(1). $\square$

Proposition 5.2.3. *Maintain the notation and assumptions of Corollary 5.2.2. The cell modules of the algebra A satisfy*

$$W(i, p) \cong \Delta_{W_i^0(p)} = A \otimes_B W_i^0(p),$$

with $W_i^0(p)$ the cell module for He_i^* , with $p \in L_i$.

Proof. This follows from Lemma 3.2.8 and Corollary 5.1.8. $\square$

5.3. Schur algebras. The following observation is implicit in [HHKP, Sections 11-13].

Lemma 5.3.1. *Consider a standardly based algebra A . The cell modules $\{W(p), p \in L\}$, form a standard system for $(L_A, \leq)$ in $A\text{-mod}$ if and only if A admits a Schur algebra as in Definition 2.9.4.*

Proof. The existence of a Schur algebra clearly implies that the cell modules form a standard system. Now consider a standardly based algebra such that $\{W(p)\}$ forms a standard system.

Set $W = \bigoplus_p W(p)$. By [DR2, Section 3], $\mathcal{F}(W)$ contains unique indecomposable objects $Y(p)$, with $p \in L$, such that $\text{Ext}_A^1(Y(p), W) = 0$ and there is a surjection $Y(p) \rightarrow W(p)$ with kernel in $\mathcal{F}(W)$. In particular $Y(\lambda) \cong P(\lambda)$ for $\lambda \in \Lambda_A \subset L_A$. Hence, we can take $m_p \in \mathbb{N}$ such that

$$A \cong \text{End}_A\left(\bigoplus_{\lambda \in \Lambda_A} P(\lambda)^{\oplus m_\lambda}\right)^{\text{op}}, \quad \text{and we define} \quad C := \text{End}_A\left(\bigoplus_{p \in L_A} Y(p)^{\oplus m_p}\right)^{\text{op}}.$$

Then C is a cover by construction and quasi-hereditary by [DR2, Section 3]. As homomorphisms and first extensions in $A\text{-mod}$ between objects in $\mathcal{F}(W)$ are categorical properties of $\mathcal{F}(W)$, this is a 1-cover by [HHKP, Lemma 11.3] or [DR2, Section 3]. $\square$

Part II. Examples

6. AUSLANDER-DLAB-RINGEL ALGEBRAS

In [Au], Auslander proved that each finite dimensional unital algebra R has a cover $A(R)$ with finite global dimension. In [DR1], Dlab and Ringel proved that $A(R)$ is quasi-hereditary. In this section, we prove that $A(R)$ admits an exact Borel subalgebra (while giving an independent proof of the quasi-heredity), under the weak assumption that R is Wedderburn.

6.1. The Auslander construction. There are several Morita equivalent versions of the algebra $A(R)$ which appear in the literature. Let R be a $\mathbb{k}$ -algebra. Consider the *right* regular R -module and its radical, which is the Jacobson radical $\mathcal{J}$ of R . For each $i \geq 1$, we let M_R^i be the largest direct summand of the *right* R -module $R/\mathcal{J}^i$ such that no direct summand of M_R^i appears as a direct summand of some M_R^j for $j < i$. We set n to be the largest integer for which $M_R^n \neq 0$. Alternatively, n is defined as the smallest positive integer for which $\mathcal{J}^n = 0$.

We define the *ADR-algebra* $A(R)$ as

$$A(R) := \text{End}_R(\oplus_{i=1}^n M_R^i).$$

We let $e_i^* \in A(R)$ be the identity morphism of M_R^i for $1 \leq i \leq n$.

We label the simple *right* R -modules by $\Lambda^0 := \Lambda_{R^{\text{op}}}$. We have unique idempotents $\{c_\lambda^* \mid \lambda \in \Lambda^0\}$ in R such that the top of the right R -module $c_\lambda^* R$ is of type λ and such that $1_R = \sum_\lambda c_\lambda^*$.

For each $\lambda \in \Lambda^0$, let $l(\lambda)$ be the Loewy length of $c_\lambda^* R$. For $l \in \mathbb{N}$ we set

$$c_{\geq l}^* = \sum_{\lambda \in \Lambda^0 \mid l(\lambda) \geq l} c_\lambda^*,$$

so in particular $c_{\geq 1}^* = 1_R$. Then we have

$$M_R^i = \bigoplus_{\lambda \in \Lambda^0 \mid l(\lambda) \geq i} c_\lambda^* R / \mathcal{J}^i = c_{\geq i}^* R / \mathcal{J}^i.$$

In particular, every direct summand of M_R^i has Loewy length i .

6.2. Construction of the Borelic subalgebra. From now on, we assume that R is Wedderburn, meaning $R = S \oplus \mathcal{J}$, for a semisimple subalgebra S . Then we have a Cartesian product of simple algebras

$$S = \bigoplus_{\lambda \in \Lambda^0} S c_\lambda^*,$$

where c_λ^* is a central idempotent in S . The space $c_\lambda^* S$ is then naturally a subspace of $c_\lambda^* R / \mathcal{J}^i$, which is bijectively mapped to the top of $c_\lambda^* R / \mathcal{J}^i$ under the canonical surjection.

We define a subspace B of $A(R)$, by

$$e_i^* B e_j^* := \{\alpha \in \text{Hom}_R(M_R^j, M_R^i) \mid \alpha(c_{\geq j}^*) \in c_{\geq i}^* S\}, \quad \forall 1 \leq i, j \leq n.$$

In particular, we find $e_i^* B e_j^* = 0$ if $i > j$, by considering Loewy lengths. If $i \leq j$, then $c_{\geq j}^* c_{\geq i}^* = c_{\geq j}^*$, so $\alpha \in e_i^* B e_j^*$ even implies $\alpha(c_{\geq j}^*) \in c_{\geq j}^* S$.

Lemma 6.2.1. *The subspace B is a graded pre-Borelic subalgebra of $A(R)$ regarded as an idempotent graded algebra.*

Proof. Set $A := A(R)$. By construction, B is a subspace of A containing 1_A . Now we prove that B is in fact an algebra. Take $\alpha \in e_i^* B e_k^*$ and $\beta \in e_k^* B e_j^*$ with $i \leq k \leq j$. It follows easily that $\alpha\beta \in B$ from the fact that S is a subalgebra of R .

To prove that B is complete, it suffices to prove equation (3.10), *viz.* that for $i \leq j$,

$$(6.1) \quad e_i^* A e_j^* = e_i^* B e_j^* \oplus \sum_{k < i} e_i^* A e_k^* A e_j^*.$$

For this we observe that the morphisms in $e_i^* B e_j^*$ are spanned by the surjections from indecomposable summands in M_R^j onto their factor modules appearing in M_R^i . A natural complement of that space is given by morphisms from M_R^j with image in the radical of M_R^i . As this image will have Loewy length strictly lower than i , the morphism factors through M_R^k (a direct sum of all factor modules of Loewy length k of the projective R -modules) for some $k < i$. This complement is hence precisely $\sum_{k < i} e_i^* A e_k^* A e_j^*$.

The algebra

$$(6.2) \quad H = B_0 \cong \bigoplus_{i=1}^n c_{\geq i}^* S$$

is semisimple and therefore ${}_H B$ is projective. Now we prove that A_B is projective. Take an $\alpha \in e_k^* A e_l^*$ which vanishes on the complement of an indecomposable direct summand N of M_R^l . Then there is an idempotent e in B such that $\alpha e = \alpha$. As argued in the previous paragraph, any morphism in eB surjects onto N , hence we find $\alpha B \cong eB$. It thus follows that A_B has a filtration with sections of the form αB , with α as above, and is therefore projective. According to Definition 3.2.3, B is a graded pre-Borelic algebra. $\square$

6.3. Main theorem on Auslander-Dlab-Ringel algebras.

Theorem 6.3.1. *Consider a field $\mathbb{k}$. If the algebra R is Wedderburn (for instance if $\mathbb{k}$ is perfect), the simple modules of the ADR algebra $A(R)$ are labelled by*

$$(6.3) \quad \Lambda_{A(R)} = \{(i, \lambda) \text{ with } 1 \leq i \leq l(\lambda) \text{ and } \lambda \in \Lambda^0\}.$$

Moreover, $(A(R), \leq_Q)$ is quasi-hereditary and admits an exact Borel subalgebra.

Proof. Since H in equation (6.2) is semisimple, the statement follows from Lemma 6.2.1, by using Corollary 4.5.4(3) and equation (3.2). $\square$

Remark 6.3.2. By [DR1], the ADR algebra is quasi-hereditary, regardless of whether R is Wedderburn. However, if R is not Wedderburn, Theorem 6.3.1 does not yield an exact Borel subalgebra. In this case, the field $\mathbb{k}$ cannot be algebraically closed and the results in [KKO] also do not ensure the existence of exact Borel subalgebras in Morita equivalence classes.

6.4. All algebras are standardly based.

Theorem 6.4.1. *Any algebra over an algebraically closed field $\mathbb{k}$ is standardly based.*

Proof. Consider an arbitrary algebra R . As the field is perfect, Theorem 6.3.1 implies that $A(R)$ is quasi-hereditary. By [DuR, Theorem 4.2.7], $A(R)$ is standardly based. As $R \cong e^* A(R) e^*$ for some idempotent e^* , it follows that R is also standardly based, by [YL, Proposition 3.5]. $\square$

6.4.2. The standardly based structure can be derived explicitly from the above proof. Take a simple module L in the socle of the right regular R -module. Acting on this with the left R -action leads to a two-sided ideal $W \otimes L$ in R for some left R -module W . Factoring out this ideal and continuing the procedure yields the structure.

Remark 6.4.3. Jie Du informed us that he was aware of this result. In the unpublished manuscript [DuL], it is proved that any split finite dimensional algebra over any field is standardly based, using the same approach as in 6.4.2.

7. EXAMPLE FROM LIE THEORY

We use the general theory to construct a very elementary Lie theoretic example of a properly stratified algebra with exact Borel subalgebra. A more advanced example can be found in [KM, Section 7]. In this section we set $\mathbb{k} = \mathbb{C}$.

7.1. **Thick category $\mathcal{O}$.** We consider the category $\mathcal{O}^k$, studied in e.g. [So]. For a reductive Lie algebra $\mathfrak{g}$, with triangular decomposition $\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$ and Borel subalgebra $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}^+$, the category $\mathcal{O}^k$ is the category of all $\mathfrak{g}$ -modules, which

- are finitely generated;
- have a basis, where each basis element v satisfies $h^k v \in \text{Span}\{h^j v \mid j < k\}$, if $h \in \mathfrak{h}$;
- are locally $U(\mathfrak{n}^+)$ -finite.

For a module M in $\mathcal{O}^k$ and $\nu \in \mathfrak{h}^*$, we consider the generalised weight space

$$M_{(\nu)} := \{v \in M \mid (h - \nu(h))^k v = 0, \text{ for all } h \in \mathfrak{h}^*\}.$$

The category $\mathcal{O}^k$ decomposes into subcategories $\mathcal{O}_\chi^k$ based on the central characters χ of $U(\mathfrak{g})$. The simple objects in $\mathcal{O}_\chi^k$ are the simple highest weight modules $L(\mu)$ with μ in the Weyl group orbit corresponding to χ , see [Hu, Section 1.7], which we denote by Λ . We consider the Bruhat (partial) order $\preceq$ on Λ of [Hu, Section 5.2]. The module $\widetilde{M}_{n,k}(\lambda)$ with $\lambda \in \Lambda$, for $n \in \mathbb{N}$, introduced in [CM, Section 4], is projective and does not depend on n , for n large enough, by [CM, Proposition 12]. We denote this module by P_λ . By the construction in [CM, Section 4], this module is generated by a vector $v_\lambda \in (P_\lambda)_{(\lambda)}$, and we have an isomorphism

$$\text{Hom}_{\mathcal{O}^k}(P_\lambda, M) \xrightarrow{\sim} M_{(\lambda)}; \quad \alpha \mapsto \alpha(v_\lambda).$$

As $\oplus_\lambda P_\lambda$ is a projective generator of $\mathcal{O}_\chi^k$, see [CM, KKM, So], we have

$$\mathcal{O}_\chi^k \cong A\text{-mod} \quad \text{with} \quad A := \text{End}_{\mathcal{O}_\chi^k} \left(\bigoplus_{\lambda \in \Lambda} P_\lambda \right)^{\text{op}}.$$

We denote the identity morphism of P_λ by e_λ^* . The modules P_λ are clearly not the indecomposable projective modules in $\mathcal{O}_\chi^k$, but they represent precisely the modules with corresponding notation in Lemma 3.1.5.

7.2. **Main theorem on thick $\mathcal{O}$.** The following theorem generalises [Kö, Theorem D].

Theorem 7.2.1. *The algebra $(A, \preceq)$ is properly stratified with exact Borel subalgebra.*

The fact that $(A, \preceq)$ is properly stratified was already pointed out in [KKM, Corollary 9(a)], due to the Morita equivalence in [So, Proposition 1].

Proof. Let us define subalgebras $B, H, \overline{N}$ of A . For arbitrary $\lambda, \mu \in \mathfrak{h}^*$, we set

$$\begin{aligned} e_\lambda^* B e_\mu^* &:= \{\alpha \in \text{Hom}_{\mathcal{O}^k}(P_\lambda, P_\mu) \mid \alpha(v_\lambda) \in U(\mathfrak{b})v_\mu\} \\ e_\lambda^* H e_\mu^* &:= \{\alpha \in \text{Hom}_{\mathcal{O}^k}(P_\lambda, P_\mu) \mid \alpha(v_\lambda) \in U(\mathfrak{h})v_\mu\} \\ e_\lambda^* \overline{N} e_\mu^* &:= \{\alpha \in \text{Hom}_{\mathcal{O}^k}(P_\lambda, P_\mu) \mid \alpha(v_\lambda) \in U(\mathfrak{n}^-)v_\mu\}. \end{aligned}$$

By construction, $e_\lambda^* H e_\mu^* = 0$ unless $\mu = \lambda$ and $H e_\lambda^* \cong \mathbb{C}[x]/(x^k)$ is a local algebra. Using the PBW theorem in [Hu, Section 0.5] and the structure of the modules $\{P_\lambda \mid \lambda \in \Lambda\}$, it can be derived that any P_μ has a basis of vectors of the form $u_1 u_2 v_\mu$, where $u_2 \in U(\mathfrak{b})$ such that $u_2 v_\mu \in (P_\mu)_{(\nu)}$ for some $\nu \in \Lambda$ and $u_1 \in U(\mathfrak{n}^-)$. This implies that $A = \overline{N}B$. This, in combination with the fact that the modules P_μ are $U(\mathfrak{n}^-)$ -free, allows one to conclude that A_B is projective. Similarly it follows that ${}_H B$ is projective. From the PBW theorem we find

$$(7.1) \quad U(\mathfrak{g}) = U(\mathfrak{n}_-)U(\mathfrak{b}) = U(\mathfrak{b}) \oplus \mathfrak{n}^-U(\mathfrak{g}) = U(\mathfrak{b}) \oplus U(\mathfrak{g})_{<}U(\mathfrak{g}),$$

where $U(\mathfrak{g})_{<}$ is the subspace of $U(\mathfrak{g})$ of all elements which are negative weight vectors for the adjoint $\mathfrak{h}$ -action. Now consider an arbitrary extension $\preceq^e$ of $\preceq$, which is a total order, meaning we can identify $(\Lambda, \preceq^e)$ with a subset of $\mathbb{N}$. The basis of P_μ mentioned above and equation (7.1) imply equation (3.10). We thus find that B is a graded exact Borelic subalgebra. The results then follow from Corollary 4.5.4(2). $\square$

8. ALGEBRAS IN DIAGRAM AND PARTITION CATEGORIES

We will obtain many examples of the types of algebras introduced in Part I, as algebras of morphisms in the partition category. This category is a natural generalisation of the Brauer category of [De, LZ] and inspired by the partition algebra of [Jo, Ma1].

8.1. Category algebras. For a category $\mathcal{C}$ with finitely many objects and morphisms, the *category algebra* $\mathbb{k}[\mathcal{C}]$ is given, as a vector space, by all formal sums of morphisms in $\mathcal{C}$ with coefficients in $\mathbb{k}$. This is an algebra for the natural product of composition. In particular, for a finite group G , which is a category with one object where all morphisms are isomorphisms, we denote the group algebra by $\mathbb{k}G$. If $\mathcal{C}$ is $\mathbb{k}$ -linear with finitely many objects and finite dimensional morphism spaces, we define the category algebra as

$$\mathbb{k}[\mathcal{C}] := \bigoplus_{x, y \in \text{Ob}\mathcal{C}} \text{Hom}_{\mathcal{C}}(x, y) \quad \text{with} \quad e_y^* \mathbb{k}[\mathcal{C}] e_x^* = \text{Hom}_{\mathcal{C}}(x, y),$$

where e_z^* is the identity morphism of $z \in \text{Ob}\mathcal{C}$.

8.2. Categories of diagrams. Consider an arbitrary field $\mathbb{k}$ and a fixed element $\delta \in \mathbb{k}$.

8.2.1. Partition category. The partition category $\mathcal{P}(\delta)$ is $\mathbb{k}$ -linear with set of objects $\mathbb{N}$. A $\mathbb{k}$ -basis of $\text{Hom}_{\mathcal{P}(\delta)}(i, j)$ is given by all partitions of the set $\{1, 2, \dots, i, 1', 2', \dots, j'\}$, which we will also view as an equivalence relation on the set.

For a partition p of the set S and a partition q of the set T , with

$$S = \{1, 2, \dots, i, 1', 2', \dots, j'\} \quad \text{and} \quad T = \{1', 2', \dots, j', 1'', 2'', \dots, k''\},$$

we define the partition $q * p$ of the set

$$S \cup T = \{1, 2, \dots, i, 1', 2', \dots, j', 1'', 2'', \dots, k''\},$$

corresponding to the minimal equivalence relation generated by p and q .

We derive two properties from $q * p$. Firstly, it induces a partition $q \odot p$ of the set

$$\{1, 2, \dots, i, 1'', 2'', \dots, k''\},$$

where two elements in the latter set are equivalent if and only if they were so in $S \cup T$. The second property is the number $d(p, q)$ of equivalence classes in $q * p$ which are contained in $S \cap T = \{1', 2', \dots, j'\}$. Now we identify the partitions p , q and $q \odot p$ with basis elements in respectively $\text{Hom}_{\mathcal{P}(\delta)}(i, j)$, $\text{Hom}_{\mathcal{P}(\delta)}(j, k)$ and $\text{Hom}_{\mathcal{P}(\delta)}(i, k)$. The product (composition) $qp = q \circ p$ is defined as $\delta^{d(p, q)} p \odot q$, which extends bilinearly to

$$\text{Hom}_{\mathcal{P}(\delta)}(j, k) \times \text{Hom}_{\mathcal{P}(\delta)}(i, j) \rightarrow \text{Hom}_{\mathcal{P}(\delta)}(i, k).$$

It is easily checked that for the above definitions, $\mathcal{P}(\delta)$ is indeed a ($\mathbb{k}$ -linear) category. For any $n \in \mathbb{Z}_{>1}$, the partition algebra is

$$\mathcal{P}_n(\delta) = \text{End}_{\mathcal{P}(\delta)}(n).$$

We will think graphically of the sets (and their partitions) by imagining the elements of $\{1, 2, \dots, i\}$ to be i dots on a horizontal line, ordered from left to right and the elements of $\{1', 2', \dots, j'\}$ to be similarly ordered on a horizontal line above the other one.

8.2.2. Brauer category. The Brauer category $\mathcal{B}(\delta)$, as introduced in [LZ, Definition 2.4], is a $\mathbb{k}$ -linear subcategory of $\mathcal{P}(\delta)$, with the same set of objects $\text{Ob}\mathcal{B}(\delta) = \mathbb{N}$. The morphism spaces are spanned by the partitions into subsets containing exactly two elements. Such partitions will be graphically represented as *Brauer diagrams*. A (k, l) -Brauer diagram consists of k points on a horizontal line, l points on a parallel line above the first line and $(k + l)/2$ lines, each connecting two points. Concatenating a (k, l) - and a (m, l) -Brauer diagram, by drawing the first on top of the second one, gives another Brauer diagram using the procedure of composing general partitions. For any $n \in \mathbb{Z}_{>1}$, the Brauer algebra is

$$\mathcal{B}_n(\delta) = \text{End}_{\mathcal{B}(\delta)}(n).$$

The lines in Brauer diagrams which connect the lower and upper line will be referred to *propagating lines*. Lines connecting two points on the lower line are called *caps* and lines connecting two points on the upper are *cups*.

8.2.3. Walled Brauer category. The walled Brauer algebra $\mathcal{B}_{r,s}(\delta)$ is a subalgebra of the Brauer algebra $\mathcal{B}_{r+s}(\delta)$ satisfying

$$\mathcal{B}_{r,s}(\delta) \cong \mathcal{B}_{s,r}(\delta) \quad \text{and} \quad \mathcal{B}_{n,0}(\delta) \cong \mathbb{k}\mathbb{S}_n.$$

There are several options to define a “walled Brauer category”. The most natural is as the category $\underline{\text{Rep}}_0(\text{GL}_\delta)$ of [CW, Section 3.2], see also [De], which has a tensor category structure. However, that category decomposes into blocks, and each such block is equivalent to a subcategory of the Brauer category $\mathcal{B}(\delta)$, which we realise as follows.

For any $p \in \mathbb{N}$, the subcategory ${}^p\mathcal{B}(\delta)$ of $\mathcal{B}(\delta)$ has set of objects $p + 2\mathbb{N} \subset \mathbb{N} = \text{Ob}\mathcal{B}(\delta)$ and the morphisms $\text{Hom}_{{}^p\mathcal{B}(\delta)}(p + 2i, p + 2j)$ are spanned by the Brauer diagrams which are “well-behaved” with respect to a straight line separating the left $p + i$ and $p + j$ dots from the right i and j dots on the two lines. Well-behaved means that propagating lines do not cross the straight line, but cups and caps intersect it precisely once. For $r \geq s > 0$, we have

$$\mathcal{B}_{r,s}(\delta) = \text{End}_{{}^{r+s}\mathcal{B}(\delta)}(r + s).$$

8.2.4. *Jones category.* A (k, l) -Jones diagram is a partition of a set of $k + l$ dots into pairs (so a Brauer diagram), which can be drawn without intersections when the l dots are on the outer boundary of an annulus and the k dots on the inner boundary of the annulus. The Jones category $J(\delta)$ is the $\mathbb{k}$ -linear subcategory of $\mathcal{B}(\delta)$, with the same set of objects $\mathbb{N}$ and where the morphisms are spanned by the Jones diagrams. For $n \in \mathbb{Z}_{>1}$, the Jones algebra is

$$J_n(\delta) = \text{End}_{J(\delta)}(n).$$

For instance, we have $J_2(\delta) \cong \mathcal{B}_2(\delta)$.

8.2.5. *Temperley-Lieb category.* The Temperley-Lieb category $\text{TL}(\delta)$ is a subcategory of the Brauer (or Jones) category with the same set of objects $\mathbb{N}$, but morphisms are spanned by the diagrams without intersections. This is the category of [BFK, Section 2.2], specialised at q with $-q - q^{-1} = \delta$. For $n \in \mathbb{Z}_{>1}$, the Temperley-Lieb algebra (of type A) is

$$\text{TL}_n(\delta) = \text{End}_{\text{TL}(\delta)}(n).$$

For instance, we have $\text{TL}_2(\delta) \cong \mathcal{B}_{1,1}(\delta)$.

8.2.6. *The category FI.* Usually, FI is introduced as the category of which the objects are all finite sets and morphisms are all injective maps between sets. We take the equivalent full subcategory with $\mathbb{N}$ as set of objects, where $n \in \mathbb{N}$ is identified with some set of cardinality n . This is a subcategory of $\mathcal{P}(\delta)$, for arbitrary δ , but is not $\mathbb{k}$ -linear, only pre-additive.

8.3. **Triangular structure and truncation.** We will distinguish three types of partitions.

- (1) Partitions of type H : These are partitions into subsets of exactly two elements, one on each line (one primed and one unprimed).
- (2) Partitions of type N^+ : These are partitions where
 - each element of the upper line is contained in a set with at least one element of the lower line and no other elements of the upper line.
 - for i , resp. j , minimal in the set containing i' , resp. j' , we have $i' < j' \Rightarrow i < j$
- (3) Partitions of type N^- : These are partitions where
 - each element of the lower line is contained in a set with at least one element of the upper line and no other elements of the lower line.
 - for i' , resp. j' , minimal in the set containing i , resp. j , we have $i < j \Rightarrow i' < j'$

By definition, partitions of type H must appear in $\text{Hom}_{\mathcal{P}(\delta)}(j, j)$, for some $j \in \mathbb{N}$, while those of type N^+ , resp. N^- , must appear in $\text{Hom}_{\mathcal{P}(\delta)}(i, j)$, for $i < j$, resp. $i > j$.

As a special case we have Brauer diagrams of the three corresponding types.

- (1) Brauer diagrams of type H : These diagrams consist solely of propagating lines.
- (2) Brauer diagrams of type N^+ : After removing all caps one is left with only non-crossing propagating lines.
- (3) Brauer diagrams of type N^- : After removing all cups one is left with only non-crossing propagating lines.

The subspace of one of the category algebras spanned by all diagrams of type H is also denoted by H , with similar convention for N^+ and N^- . For each of the categories, except ${}^p\mathcal{B}(\delta)$, we denote the identity morphism in $\text{End}(i)$ by e_i^* . For ${}^p\mathcal{B}(\delta)$ we use e_i^* for the identity morphism of $p + 2i \in \text{Ob}{}^p\mathcal{B}(\delta)$.

For each $n \in \mathbb{Z}_{>1}$, we introduce the full subcategory $\mathcal{P}^{\leq n}(\delta)$, resp. $\text{FI}^{\leq n}$, of $\mathcal{P}(\delta)$, resp. FI, with objects $\{0, 1, 2, \dots, n\} \subset \mathbb{N}$. Their category algebras satisfy (3.7).

We observe that the diagram categories always decompose into subcategories with as objects either even or odd natural numbers. Hence, for $n \in \mathbb{Z}_{>1}$, we define $\mathcal{B}^{\leq n}(\delta)$, $J^{\leq n}(\delta)$ and $\text{TL}^{\leq n}(\delta)$ as the full subcategories of resp. $\mathcal{B}(\delta)$, $J(\delta)$ and $\text{TL}(\delta)$, with objects

$$(8.1) \quad \mathcal{J}(n) := \{n, n-2, \dots, n-2\lfloor \frac{n}{2} \rfloor\}.$$

The corresponding category algebras satisfy $1 = \sum_{i \in \mathcal{J}(n)} e_i^*$. For $p \in \mathbb{N}$ and $n \in \mathbb{Z}_{>0}$, the category ${}^p\mathcal{B}^{\leq n}(\delta)$ is the full subcategory of ${}^p\mathcal{B}(\delta)$ with objects

$$\{p, p+2, \dots, p+2n\}.$$

The category algebra satisfies equation (3.7).

8.4. The category algebras. We consider the category algebras of the previous section with idempotents e_i^* . We recall the corresponding decomposition $\mathcal{Q}$ of Λ , and the idempotent quasi-order $\preceq_{\mathcal{Q}}$ and idempotent partial order $\leq_{\mathcal{Q}}$ on Λ of Definition 4.5.1.

Theorem 8.4.1. *Fix an arbitrary field $\mathbb{k}$ and $\delta \in \mathbb{k}$.*

- (1) *The algebras $\mathbb{k}[\mathcal{P}^{\leq n}(\delta)]$, $\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$, $\mathbb{k}[{}^p\mathcal{B}^{\leq n-p}(\delta)]$ and $\mathbb{k}[\text{FI}^{\leq n}(\delta)]$ are*
 - *quasi-hereditary for $\leq_{\mathcal{Q}}$, with exact Borel subalgebra if $\text{char}(\mathbb{k}) \notin [2, n]$;*
 - *exactly standardly stratified for $\preceq_{\mathcal{Q}}$, with exact Borel subalgebra;*
 - *base stratified for decomposition $\mathcal{Q}$.*
- (2) *The algebra $\mathbb{k}[J^{\leq n}(\delta)]$ is*
 - *quasi-hereditary for $\leq_{\mathcal{Q}}$ with exact Borel subalgebra, if $\text{char}(\mathbb{k})$ does not divide any element of $\mathcal{J}(n)$;*
 - *properly stratified for $\leq_{\mathcal{Q}}$, with exact Borel subalgebra;*
 - *base stratified for decomposition $\mathcal{Q}$ if the polynomials $x^i - 1$, for $i \in \mathcal{J}(n)$, split over the field $\mathbb{k}$.*
- (3) *The algebra $\mathbb{k}[\text{TL}^{\leq n}(\delta)]$ is*
 - *quasi-hereditary for $\leq_{\mathcal{Q}}$, with exact Borel subalgebra;*
 - *base stratified for decomposition $\mathcal{Q}$.*

When the above condition for quasi-heredity is not satisfied, the category algebra is not quasi-hereditary, for any partial order on Λ (with Λ given in Lemma 8.4.5).

Remark 8.4.2. The statements on the exact standard stratification in the theorem can be refined by replacing $\preceq_{\mathcal{Q}}$ by a partial quasi-order $\preceq$, such that $<$ is the same as $\prec_{\mathcal{Q}}$, but $\sim$ reflects the block decomposition of $\mathbb{k}\mathbb{S}_t$, see [Ja, Section 6.1].

If $\mathbb{k}$ is algebraically closed, the algebras in Theorem 8.4.1 are all standardly based by Theorem 5.1.7, with poset $L := \sqcup_i L_i$ and L_i the posets for the standardly based algebra He_i^* , as can be obtained from Examples 2.9.2 and 2.9.3 and Lemma 8.4.5.

Theorem 8.4.3. *Assume that $\mathbb{k}$ is algebraically closed. The cell modules of C form a standard system for $(L, \leq)$ if and only if the condition given below is satisfied.*

algebra C	condition	Set $L = \sqcup_i L_i$
$\mathbb{k}[\mathcal{P}^{\leq n}(\delta)]$	$\text{char}(\mathbb{k}) \notin \{2, 3\}$ or $\text{char}(\mathbb{k}) = 3$ and $n = 2$	$\{(i, \mu) \mid 0 \leq i \leq n, \mu \vdash i\}$
$\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$	$\text{char}(\mathbb{k}) \notin \{2, 3\}$ or $\text{char}(\mathbb{k}) = 3$ and $n = 2$	$\{(i, \mu) \mid i \in \mathcal{J}(n), \mu \vdash i\}$
$\mathbb{k}[J^{\leq n}(\delta)]$	i not divisible by $\text{char}(\mathbb{k})$, for $i \in \mathcal{J}(n)$	$\{(i, \omega) \mid i \in \mathcal{J}(n), \omega \in \mathbb{k}, \omega^i = 1\}$
$\mathbb{k}[\text{TL}^{\leq n}(\delta)]$	$\emptyset$	$\mathcal{J}(n)$
$\mathbb{k}[\text{FI}^{\leq n}(\delta)]$	$\text{char}(\mathbb{k}) \notin \{2, 3\}$ or $\text{char}(\mathbb{k}) = 3$ and $n = 2$	$\{(i, \mu) \mid 0 \leq i \leq n, \mu \vdash i\}$
$\mathbb{k}[{}^p\mathcal{B}^{\leq n}(\delta)]$	$\text{char}(\mathbb{k}) \notin \{2, 3\}$ or $\text{char}(\mathbb{k}) = 3$ and $n + p \leq 2$	$\{(i, \mu, \nu) \mid 0 \leq i \leq n, \mu \vdash p + i, \nu \vdash i\}$

We start the proofs of the theorems with the following proposition.

Proposition 8.4.4. *Let C be one of the category algebras in Theorem 8.4.3, equipped with the idempotent grading of (3.8). The subspaces H, N^+ and N^- of C , defined in Section 8.3 are subalgebras and $B := HN^+$ is a graded pre-Borelic subalgebra of C .*

Proof. One verifies that the subspaces are subalgebras for each case. Further, we can classify partitions into those where each element of the upper line is contained in a set/class with at least one element of the lower line and no other elements of the upper line, and the rest. In each of the cases, the first type of diagrams span B , whereas those of the second type span C_-B , proving equation (3.9). So we find that B is a complete $\mathbb{Z}_+$ -graded subalgebra of C .

It hence remains to prove that C_B and ${}_HB$ are projective modules. We have $C = N^-B$ and by associating to each partition the number of classes which contain elements of both rows (for diagrams this is just the number of propagating lines) we find a decomposition

$$C = \bigoplus_i N^- e_i^* B.$$

For any partition q in $N^- e_i^*$ we have $qB \cong e_i^* B$, proving that C_B is projective. Similarly it follows that for any partition q in $e_i^* B$ we have $Hq \cong He_i^*$, which concludes the proof. $\square$

For the cyclic group C_t of order t , we have $\mathbb{k}C_t \cong \mathbb{k}[x]/(x^t - 1)$. We denote its labelling set of simple modules by $\Lambda_C^{\mathbb{k}, t}$. In case the polynomial $x^t - 1$ splits over $\mathbb{k}$, for instance when $\mathbb{k}$ is algebraically closed, we can take $\{\omega \in \mathbb{k} \mid \omega^t = 1\}$ for this set.

Lemma 8.4.5. *Let C be any of the category algebras in Proposition 8.4.4. The following table summarises the structure of the subalgebra $H \subset C$, its labelling set Λ_C of isoclasses of simple modules and the criterion for H to be semisimple.*

algebra C	algebra H	set $\Lambda_C = \Lambda_H = \sqcup_i \Lambda_i$	semisimplicity criterion for H
$\mathbb{k}[\mathcal{P}^{\leq n}(\delta)]$	$\bigoplus_{i=0}^n \mathbb{k}\mathcal{S}_i$	$\{(i, \mu) \mid 0 \leq i \leq n, \mu \vdash_{\mathbb{k}} i\}$	$\text{char}(\mathbb{k}) \notin [2, n]$
$\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$	$\bigoplus_{i \in \mathcal{J}(n)} \mathbb{k}\mathcal{S}_i$	$\{(i, \mu) \mid i \in \mathcal{J}(n), \mu \vdash_{\mathbb{k}} i\}$	$\text{char}(\mathbb{k}) \notin [2, n]$
$\mathbb{k}[J^{\leq n}(\delta)]$	$\bigoplus_{i \in \mathcal{J}(n)} \mathbb{k}C_i$	$\{(i, \omega) \mid i \in \mathcal{J}(n), \omega \in \Lambda_C^{\mathbb{k}, i}\}$	i not divisible by $\text{char}(\mathbb{k})$, $i \in \mathcal{J}(n)$
$\mathbb{k}[\text{TL}^{\leq n}(\delta)]$	$\bigoplus_{i \in \mathcal{J}(n)} \mathbb{k}$	$\mathcal{J}(n)$	$\emptyset$
$\mathbb{k}[\text{FI}^{\leq n}(\delta)]$	$\bigoplus_{i=0}^n \mathbb{k}\mathcal{S}_i$	$\{(i, \mu) \mid 0 \leq i \leq n, \mu \vdash_{\mathbb{k}} i\}$	$\text{char}(\mathbb{k}) \notin [2, n]$
$\mathbb{k}[{}^p\mathcal{B}^{\leq n}(\delta)]$	$\bigoplus_{i=0}^n \mathbb{k}\mathcal{S}_{p+i}$ $\oplus \bigoplus_{i=0}^n \mathbb{k}\mathcal{S}_i$	$\{(i, \mu, \nu) \mid 0 \leq i \leq n, \mu \vdash_{\mathbb{k}} p + i, \nu \vdash_{\mathbb{k}} i\}$	$\text{char}(\mathbb{k}) \notin [2, p + n]$

Proof. The structure of the algebra H follows from definition in Section 8.2. We use equation (3.2). The simple modules of $\mathbb{k}\mathcal{S}_t$ are naturally labelled by $\mathbb{k}$ -regular partitions of t , see

[Ja, Section 6.1]. By Maschke's theorem, for a finite group G , the algebra $\mathbb{k}G$ is semisimple if and only if the order $|G|$ is not divisible by $\text{char}(\mathbb{k})$. Hence, $\mathbb{k}\mathbb{S}_t$ is semisimple if and only if $\text{char}(\mathbb{k}) > t$ or $\text{char}(\mathbb{k}) = 0$. $\square$

Proof of Theorem 8.4.1. We freely use Proposition 8.4.4 and Lemma 8.4.5.

That the algebras are based-stratified follows from Theorem 5.2.1 and Examples 2.9.2 and 2.9.3. The exact standard stratification follows from Corollary 4.5.4(1). The quasi-heredity follows from Corollary 4.5.4(3). It is easily checked that the group algebra $\mathbb{k}C_i$ is quasi-local, hence $C = \mathbb{k}[J^{\leq n}(\delta)]$ is properly stratified by Corollary 4.5.4(2).

Now we prove that in the remaining cases C is not quasi-hereditary. By equation (3.14) and Lemma A.1.1, C will have infinite global dimension as soon as H has. For any finite group G , the algebra $\mathbb{k}G$ is Frobenius and hence self-injective. In particular, the global dimension of $\mathbb{k}G$ is finite if and only if it is zero. In conclusion, when the criteria for semisimplicity of H in Lemma 8.4.5 are not satisfied, C has infinite global dimension. In particular C is not quasi-hereditary for any order, by [CPS1] or [APT, Corollary 6.6]. $\square$

The following lemma can be derived from [KN] with minor additional work.

Lemma 8.4.6. *Assume that $\mathbb{k}$ is algebraically closed and $i > 1$.*

- *The cell modules of the standardly based algebra $\mathbb{k}\mathbb{S}_i$ of Example 2.9.3 form a standard system if and only if*

$$\begin{cases} \text{char}(\mathbb{k}) \notin \{2, 3\} & \text{or} \\ \text{char}(\mathbb{k}) = 3 & \text{and } i = 2. \end{cases}$$

- *The cell modules of the standardly based algebra $\mathbb{k}C_i$ of Example 2.9.2 form a standard system if and only if $\text{char}(\mathbb{k})$ does not divide i .*

Proof. For $\text{char}(\mathbb{k}) \notin \{2, 3\}$, [KN, Theorem 6.4(b)(ii)] implies that the Specht modules of $\mathbb{k}\mathbb{S}_i$ form a standard system. When $\text{char}(\mathbb{k}) = 3$, $\mathbb{k}\mathbb{S}_2$ is semisimple and the (simple) Specht modules form a standard system.

Now we prove that, in the remaining cases, the Specht modules do not constitute a standard system, for any order. The trivial module S_1 and the sign module S_2 are both Specht modules. Assume $\text{char}(\mathbb{k}) = 2$, then $S_1 \cong S_2$, contradicting property (2) in the definition of a standard system in 2.7. Now assume $\text{char}(\mathbb{k}) = 3$ and $i \geq 3$. We will prove that

$$\text{Ext}_{\mathbb{k}\mathbb{S}_i}^1(S_1, S_2) \neq 0 \neq \text{Ext}_{\mathbb{k}\mathbb{S}_i}^1(S_2, S_1),$$

contradicting property (3) in the definition of a standard system. The modules M and N corresponding to the extensions are given by

$$M = \langle v, w \rangle \quad \text{with} \quad s_j v = v + c_j w \quad \text{and} \quad s_j w = -w,$$

for $s_j, j \in \{1, \dots, i-1\}$ the generators of $\mathbb{S}_i$ and $c_j \in \mathbb{k}$ arbitrary, and

$$N = \langle v, w \rangle \quad \text{with} \quad s_j w = -w + d_j v \quad \text{and} \quad s_j v = v,$$

for $d_j \in \mathbb{k}$ arbitrary. It can be verified that in characteristic 3, there are no conditions on the coefficients c_j, d_j coming from imposing the braid relations. However, when $c_j \neq c_{j'}$ or $d_j \neq d_{j'}$ the modules do not decompose and the extensions hence do not split.

For $\mathbb{k}C_i \cong \mathbb{k}[x]/(x^i - 1)$, the cell modules are two-by-two non-isomorphic if and only if all i -th roots of 1 in the algebraically closed field $\mathbb{k}$ are different. This is equivalent to demanding

that $\mathbb{k}C_i$ is semisimple, in which case the cell modules are simple and form a standard system for any order. So a necessary and sufficient condition is that $\text{char}(\mathbb{k})$ does not divide i . $\square$

Proof of Theorem 8.4.3. By Corollary 5.2.2 and Proposition 8.4.4, this follows from Lemmata 8.4.5 and 8.4.6. $\square$

Remark 8.4.7. Consider the case $\text{char}(\mathbb{k}) \notin [2, n]$. We have proved that $\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$ is quasi-hereditary with exact Borel subalgebra. This is not a *strong* exact Borel subalgebra as defined in [Kö], as H is not a maximal semisimple subalgebra. In fact, generically, $\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$ will be semisimple itself, as follows from [Ru] and Theorem 8.5.1 below.

Remark 8.4.8. The exact Borel subalgebra B of $C := \mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$ does not satisfy the property

$$\text{Ext}_C^k(C \otimes_B M, C \otimes_B N) \cong \text{Ext}_B^k(M, N), \quad \forall k \geq 2,$$

for arbitrary $M, N \in C\text{-mod}$, which is satisfied for the exact Borel subalgebra predicted by the general theory of [KKO], see [KKO, Theorems 1.1 and 10.5]. We observe that B , and hence the extension group in the right-hand side, does not depend on δ . The extension group in the left-hand side depends heavily on δ . As mentioned in Remark 8.4.7, for generic δ the left-hand side will vanish. It is easily checked that it does not vanish for *e.g.* $\delta = 0$. This proves that the displayed isomorphism cannot be true in general.

Remark 8.4.9. For $C = \mathbb{k}[\text{FI}^{\leq n}]$, we have $B = H$ as $N^+ = 0$. So for $\text{char}(\mathbb{k}) \notin [2, n]$, $\mathbb{k}[\text{FI}^{\leq n}]$ is quasi-hereditary with semisimple exact Borel subalgebra. It can easily be checked that for the reversal of the order $\leq_Q$, we have that $\mathbb{k}[\text{FI}^{\leq n}]$ is an exact Borel subalgebra of itself.

8.5. Morita equivalences.

Theorem 8.5.1. *For an arbitrary field $\mathbb{k}$, we have the following Morita equivalences under the respective conditions given in the table:*

Morita equivalence	condition
$\mathcal{P}_n(\delta) \stackrel{M}{=} \mathbb{k}[\mathcal{P}^{\leq n}(\delta)]$	$\delta \neq 0$
$\mathcal{B}_n(\delta) \stackrel{M}{=} \mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$	$\delta \neq 0$ or n odd
$J_n(\delta) \stackrel{M}{=} \mathbb{k}[J^{\leq n}(\delta)]$	$\delta \neq 0$ or n odd
$\text{TL}_n(\delta) \stackrel{M}{=} \mathbb{k}[\text{TL}^{\leq n}(\delta)]$	$\delta \neq 0$ or n odd
$\mathcal{B}_{p+n,n}(\delta) \stackrel{M}{=} \mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$	$\delta \neq 0$ or $p \neq 0$

Remark 8.5.2. Theorem 8.5.1 and Lemma 8.4.5 together determine the labelling set Λ_A of the simple modules of the diagram algebras A , under the condition in the right column of the table. In almost all cases Λ_A is known by [CDDM, GL, Ma2, Xi].

We start the proof of Theorem 8.5.1 by constructing special elements in the category algebras, which will also be essential for constructions in Part III.

Lemma 8.5.3. *Consider algebras C , $A = e_n^* C e_n^*$ and $i \in I_A(n)$, as in the table below.*

Category algebra C	Diagram algebra A	$I_A(n)$
$\mathbb{k}[\mathcal{P}^{\leq n}(\delta)]$	$\mathcal{P}_n(\delta)$	$\{0, 1, \dots, n\}$
$\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$	$\mathcal{B}_n(\delta)$	$\mathcal{J}(n)$
$\mathbb{k}[J^{\leq n}(\delta)]$	$J_n(\delta)$	$\mathcal{J}(n)$
$\mathbb{k}[\text{TL}^{\leq n}(\delta)]$	$\text{TL}_n(\delta)$	$\mathcal{J}(n)$
$\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$	$\mathcal{B}_{p+n,n}(\delta)$	$\{0, 1, \dots, n\}$

Then there are elements $a_i \in e_n^* C e_i^*$ and $b_i \in e_i^* C e_n^*$ such that $c_i^* := a_i b_i \in A$ and e_i^* satisfy

$$\begin{cases} e_i^* = b_i a_i & \text{and } (c_i^*)^2 = c_i^* & \text{if either } \delta \neq 0 \text{ or } i + p \neq 0 \\ (c_i^*)^2 = 0 & & \text{if } \delta = 0 \text{ and } i + p = 0. \end{cases}$$

Here, we set $p = 0$ in all cases, except for ${}^p\mathcal{B}^{\leq n}(\delta)$.

Proof. There are many different choices for a_i and b_i . The constructions given below for the various cases produce different a_i and b_i in the overlapping situations.

Firstly we consider the partition category (so $p = 0$) and arbitrary $1 \leq i \leq n$. We put

$$b_i = \{\{1, 1'\}, \{2, 2'\}, \dots, \{i, i'\}, \{i+1, i+2, \dots, n\}\} \quad \text{and} \\ a_i = \{\{1, 1'\}, \{2, 2'\}, \dots, \{i-1, (i-1)'\}, \{i, i', (i+1)', (i+2)', \dots, n'\}\}.$$

We have $b_i a_i = e_i^*$, which implies that $c_i^* := a_i b_i$ is an idempotent. In case $i = 0$ we define $a_0 \in \text{Hom}_{\mathcal{P}(\delta)}(0, n)$ and $\bar{a}_0 \in \text{Hom}_{\mathcal{P}(\delta)}(n, 0)$ as the partitions into one set. When $\delta \neq 0$, we set $b_0 = \bar{a}_0/\delta$, which leads to $b_0 a_0 = e_0^*$ and an idempotent $c_0^* = a_0 b_0$. When $\delta = 0$ we set $b_0 = \bar{a}_0$, in which case $c_0^* := a_0 b_0$ squares to zero. The above already deals with the partition category completely, although the diagrams that will be constructed below for the other categories can also be used for the partition category when $i \in \mathcal{J}(n)$.

Now we consider the cases $\mathcal{B}_n(\delta)$, $J_n(\delta)$ and $\text{TL}_n(\delta)$ (so $p = 0$). For arbitrary $i \in \mathcal{J}(n)$ we introduce three Brauer diagrams, which are also Temperley-Lieb and hence Jones diagrams. We consider two diagrams $\bar{a}_i$ and $\hat{a}_i$ with $(n-i)/2$ caps and i propagating lines and a_i with $(n-i)/2$ cups and i propagating lines. Note that the definition of $\hat{a}_i$ requires $i > 0$.

$$\begin{aligned} \bar{a}_i &= \left| \begin{array}{c} | \quad | \quad \cdots \quad | \quad | \quad \cup \quad \cup \quad \cdots \quad \cup \end{array} \right. \\ \hat{a}_i &= \left| \begin{array}{c} | \quad | \quad \cdots \quad | \quad \cup \quad \cup \quad \cdots \quad \cup \end{array} \right. \\ a_i &= \left| \begin{array}{c} | \quad | \quad \cdots \quad | \quad | \quad \cup \quad \cup \quad \cdots \quad \cup \end{array} \right. \end{aligned}$$

If $\delta \neq 0$, we can set $b_i := \delta^{(i-n)/2} \bar{a}_i$ and then we have $b_i a_i = e_i^*$. If $i \neq 0$, we can set $b_i := \hat{a}_i$ and then we have $b_i a_i = e_i^*$. In either case, we automatically find that $c_i^* := a_i b_i$ is then an idempotent. When both $\delta = 0$ and $i = 0$, we set $b := \bar{a}_0$ and then $c_0^* = a_0 b_0$ squares to zero. This completes the cases $\mathcal{B}_n(\delta)$, $J_n(\delta)$ and $\text{TL}_n(\delta)$.

The case $\mathcal{B}_{p+n,n}(\delta)$ behaves similarly. For instance, if $p > 0$ and $\delta = 0$, we can take

$$c_0^* := \left| \begin{array}{c} | \quad | \quad \cdots \quad | \quad | \quad \cup \quad \cup \quad \cdots \quad \cup \end{array} \right.$$

where there are p propagating lines and n cups and caps. □

Proof of Theorem 8.5.1. Any of the category algebras satisfies $1_C = \sum_{i=0}^n e_i^*$. The condition in the right column of the table in the theorem is precisely that $\delta \neq 0$ or $e_0^* = 0$. Under this condition, Lemma 8.5.3 implies that $1_C \in C e_n^* C$, and hence $C = C e_n^* C$. The conclusion thus follows from case (c) of Section 2.4. □

8.6. Cellularity. We prove that the category algebras and the diagram algebras are cellular in the sense of [GL, Definition 1.1].

8.6.1. For all $i, j \in \mathbb{N}$, we define a $\mathbb{k}$ -vector space isomorphism

$$\iota : \text{Hom}_{\mathcal{P}(\delta)}(i, j) \rightarrow \text{Hom}_{\mathcal{P}(\delta)}(j, i),$$

which is determined by demanding that any partition is mapped to its “horizontal flip”, which simply identifies the i dots on the lower line of partitions in $\text{Hom}_{\mathcal{P}(\delta)}(i, j)$ with those on the upper line of partitions in $\text{Hom}_{\mathcal{P}(\delta)}(j, i)$. This clearly extends to an involutive anti-algebra morphism of $\mathbb{k}[\mathcal{P}^{\leq n}(\delta)]$, which furthermore restricts to involutive anti-algebra morphisms of $\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$, $\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$ and $\mathbb{k}[\text{TL}^{\leq n}(\delta)]$. On a diagram with only propagating lines, interpreted as an element of a symmetric group, ι acts as inversion. For $\mathbb{k}[J^{\leq n}(\delta)]$ we use a different involution ι , *viz.* ι is defined on a Jones diagram as its horizontal and vertical flip. In particular, this stabilises any Jones diagram with only propagating lines.

Theorem 8.6.2. *Consider an arbitrary field $\mathbb{k}$. Let C be $\mathbb{k}[\mathcal{P}^{\leq n}(\delta)]$, $\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$, $\mathbb{k}[J^{\leq n}(\delta)]$, $\mathbb{k}[\text{TL}^{\leq n}(\delta)]$ or $\mathbb{k}[\mathcal{B}^{\leq n}(\delta)]$. Then C is cellular for involution ι when it satisfies the condition to be base stratified in Theorem 8.4.1.*

Proof. We take an arbitrary completion of the partial order of 5.1.6 on $L = \sqcup_i L_i$, again denoted by $\leq$. The ideals $C^{\geq(l,p)}$ for $(l, p) \in L$ of the standardly based algebra C then form a chain of ideals, which is refinement of the chain of ideals $J_i = C f_i C = C^{\geq(i, p_0)}$, with p_0 minimal in L_i . We introduce the space J'_i spanned by all partitions where exactly i subsets contain dots of the upper and lower row, or the intersection of that space with the relevant category algebra. Then we have

$$J_i = J'_i \oplus J_{i-1} \quad \text{and} \quad \iota(J'_i) = J'_i.$$

By extending the cellular structures on He_i^* in Examples 2.9.2 and 2.9.3 to C , we construct similar decompositions

$$C^{\geq(l,p)} = (C^{\geq(l,p)})' \oplus C^{\geq(l,p_1)}$$

for the refinement, where p_1 is the (unique) maximal $q \in L_l$ with $q < p$. Together with the above decompositions, it then follows easily that [KX2, Definition 2.2] is satisfied for the refined chain and C is hence cellular, see also [DuR, Lemma 1.2.4]. $\square$

As $\iota(e_n^*) = e_n^*$, [KX1, Proposition 4.3] implies the following property.

Corollary 8.6.3. *The algebras $\mathcal{P}_n(\delta)$, $\mathcal{B}_n(\delta)$, $\mathcal{B}_{r,s}(\delta)$ and $\text{TL}_n(\delta)$ are cellular with respect to the restriction of ι . If the polynomials $x^i - 1$, for $i \in \mathcal{J}(n)$, split over $\mathbb{k}$, the algebra $J_n(\delta)$ is cellular with respect to the restriction of ι .*

Proposition 8.6.4. *Consider any of the cellular algebras A in Corollary 8.6.3 with the corresponding category algebra C in Lemma 8.5.3. The cell modules of A are labelled by $L_A := L_C$ in Theorem 8.4.3, and are given by*

$$W_A(i, p) := e_n^* W_C(i, p) \quad \text{with} \quad W_C(i, p) = C \otimes_B W_i^0(p) \quad \text{for } i \in I_A \text{ and } p \in L_i.$$

Proof. We use Proposition 5.2.3. Since

$$e_n^* C \otimes_B W_i^0(p) \neq 0,$$

for any $W_i^0(p)$, the proposition follows from the proof of [KX1, Proposition 4.3]. $\square$

Part III. Faithfulness of covers

9. COVERS OF THE DIAGRAM ALGEBRAS

The two main methods to prove that cell modules of a cellular algebra form a standard system are construction of a certain quasi-hereditary 1-cover (*viz.* a Schur algebra), see [KN, HK]; and cellular/base stratification, see [HHKP] and Section 8.

When the diagram algebra A in Lemma 8.5.3 is Morita equivalent to C , the base stratification of C solves the problem also for A . In this section, we use a combination of both approaches above, for when A is not Morita equivalent to C . We prove that the based stratified algebra C is almost always a 1-faithful cover of A , in the sense of Section 2.8. Even though C might not be quasi-hereditary, this determines when the cell modules of A form a standard system.

9.1. Connection with the coarse filtration. In Section 8.2, we defined the diagram algebras as the endomorphism algebras of the objects of the corresponding category. Remarkably, we can reconstruct the respective categories starting from the diagram algebra, by using the elements $\{c_i^* \mid i \in I_A(n)\}$ of Lemma 8.5.3.

Theorem 9.1.1. *Consider an arbitrary field $\mathbb{k}$, $\delta \in \mathbb{k}$ and let $\mathcal{A}$ be $\mathcal{P}(\delta)$, $\mathcal{B}(\delta)$, ${}^p\mathcal{B}(\delta)$, $J(\delta)$ or $\text{TL}(\delta)$. For $n \in \mathbb{Z}_{>0}$, set $A := \text{End}_{\mathcal{A}}(n)$. For any $i, j \in I_A(n)$, we have*

$$\text{Hom}_{\mathcal{A}}(j, i) \cong \text{Hom}_A(Ac_i^*, Ac_j^*).$$

Moreover, the composition of such morphisms on both sides agrees contravariantly.

As a special case we have the following corollary, giving an alternative description of the category algebra C connected to the diagram algebra A .

Corollary 9.1.2. *Maintain the notation of Theorem 9.1.1 and consider the category algebra $C := \mathbb{k}[A^{\leq n}]$. We have an isomorphism of left A -modules*

$$Ac_j^* \cong \text{Hom}_{\mathcal{A}}(j, n) = e_n^* C e_j^*, \quad \text{for } j \in I_A(n),$$

and an isomorphism of algebras

$$C \cong \text{End}_A\left(\bigoplus_{j \in I_A(n)} Ac_j^*\right)^{\text{op}}.$$

9.1.3. Before we prove the theorem, we elaborate on the idempotents c_i^* . The partition algebra has a filtration by two-sided ideals, known as the *coarse filtration*:

$$0 = J_0 \subsetneq J_1 \subsetneq J_2 \subsetneq \cdots \subsetneq J_n \subsetneq J_{n+1} = \mathcal{P}_n(\delta),$$

see *e.g.* [Xi, Lemma 4.6]. Here, J_i is the ideal spanned by those partitions of

$$(9.1) \quad S = S_1 \sqcup S_2 = \{1, 2, \dots, n\} \sqcup \{1', 2', \dots, n'\},$$

where at most $i-1$ of the subsets contain elements of both S_1 and S_2 , for each $1 \leq i \leq n+1$. Note that we have $J_i = Ac_{i-1}^* A$ for $1 \leq i \leq n+1$. Hence, J_i is an idempotent ideal, if either $i \neq 1$ or $\delta \neq 0$. When $\delta = 0$, we have $J_1^2 = 0$.

Remark 9.1.4. When $\delta \neq 0$, the coarse filtration is an exactly standardly stratifying chain, even quasi-heredity when $\text{char}(\mathbb{k}) \notin [2, n]$, which leads to the corresponding properties of $\mathcal{P}_n(\delta)$ in Theorem 2. Furthermore (for any $\mathbb{k}$ and $\delta \in \mathbb{k}$) the chain can be refined to a cell chain, see [KX1, Definition 3.2], expressing the cellular structure in Theorem 2.

9.1.5. By restricting the coarse filtration of $\mathcal{P}_n(\delta)$, we obtain coarse filtrations of $\mathcal{B}_n(\delta)$, $\mathcal{B}_{r,s}(\delta)$, $J_n(\delta)$ and $\text{TL}_n(\delta)$. These coarse filtrations are based on the number of propagating lines. The corresponding idempotents (or the nilpotent element) generating the ideals are also given by $\{c_i^*\}$ in Lemma 8.5.3.

Now we start the proof of Theorem 9.1.1 with the following lemma.

Lemma 9.1.6. *Let A be one of the diagram algebras in Theorem 9.1.1, with n even in case A is $\mathcal{B}_n(\delta)$, $J_n(\delta)$ or $\text{TL}_n(\delta)$. We have an isomorphism*

$$\text{Hom}_A(Ac_0^*, A) \xrightarrow{\sim} c_0^*A, \quad \alpha \mapsto \alpha(c_0^*).$$

Proof. We have a monomorphism

$$\phi : \text{Hom}_A(Ac_0^*, A) \hookrightarrow A, \quad \alpha \mapsto \alpha(c_0^*).$$

First we claim that c_0^*A is contained in the image of ϕ . We have the A -bimodule isomorphism

$$Ac_0^* \otimes_{\mathbb{k}} c_0^*A \xrightarrow{\sim} Ac_0^*A; \quad ac_0^* \otimes c_0^*b \mapsto ac_0^*b,$$

which is clear by construction, or follows from the cellular structure. In particular, for any $x \in c_0^*A$, the above gives rise to a morphism α_x from Ac_0^* to A , which maps c_0^* to x . Since $\phi(\alpha_x) = x$, this proves the claim.

Now we will prove that the image of ϕ is contained in c_0^*A . If c_0^* is an idempotent, this follows from $\alpha(c_0^*) = c_0^*\alpha(c_0^*)$. Consider therefore $(c_0^*)^2 = 0$ and some element $a \in A$ in the image of ϕ , so $a = \alpha(c_0^*)$ for some α . We prove that $a \in c_0^*A$ by considering cases separately.

Firstly, consider $A = \mathcal{P}_n(0)$. Observe that a must satisfy $c_1^*a = a$, since $c_1^*c_0^* = c_0^*$. Hence, a must be a linear combination of partitions of (9.1) where $S_2 = \{1', \dots, n'\}$ is contained in a subset. So $a = a_1 + a_2$, where a_1 is a linear combination of partitions where S_2 is a subset (hence $a_1 \in c_0^*A$) and a_2 is a linear combination of partitions which have a subset which strictly contains S_2 . We define $u = u_1 - u_2$, with u_1, u_2 partitions of (9.1) as:

$$(9.2) \quad u_1 := \{\{1, 2, \dots, n, 1'\}, \{2', 3', \dots, n'\}\}, \quad u_2 := \{\{1, 2, \dots, n, 2', 3', \dots, n'\}, \{1'\}\}.$$

Since $uc_0^* = 0 = ua_1$, we must have $ua_2 = 0$. However, u_1a_2 is a linear combination of partitions where $\{1'\} \cup S_1$ is a subset, whereas u_2a_2 is a linear combination of partitions where $S \setminus \{1'\}$ is a subset. Hence $u_1a_2 = 0 = u_2a_2$. As the horizontal flip v_1 of the partition u_1 satisfies $v_1u_1a_2 = a_2$, this shows that $a_2 = 0$ and hence that $a \in c_0^*A$.

Secondly, consider A equal to $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$ with n even. As the case $n = 2$ is straightforward, we assume $n \geq 4$. We now have that $c_2^*a = a$. This means a is a linear combination of diagrams with top row of the form

$$(9.3) \quad \cdot \quad \cdot \quad \cup \quad \cup \quad \dots \quad \cup$$

where the first two dots can be arbitrarily connected. Consider also the diagram

$$(9.4) \quad w := \begin{array}{c} \diagup \quad \diagdown \quad | \quad | \quad | \quad \dots \quad | \quad | \\ \diagdown \quad \diagup \end{array}$$

Since $wc_0^* = c_0^*$ we must also have $wa = a$. This implies that the top rows of the diagrams (9.3), appearing in a , must have only caps. It now follows easily that $a \in c_0^*A$.

Now set $A := \mathcal{B}_{n,n}(0)$ and use c_0^* and c_1^* as in 9.2.5. We focus on the case $r > 1$ and define

$$(9.5) \quad v := \begin{array}{c} \text{Diagram showing a sequence of vertical lines and curved lines, representing a diagram in the category } \mathcal{A}. \end{array}$$

which satisfies $vc_0^* = c_0^*$. The conclusion follows as for the previous case. $\square$

Lemma 9.1.7. *Maintain the notation of Theorem 9.1.1. We have an isomorphism*

$$\mathrm{Hom}_{\mathcal{A}}(j, i) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{A}}(Ac_i^*, Ac_j^*); \quad x \mapsto \alpha_x \quad \text{with} \quad \alpha_x(c_i^*) = a_i x b_j,$$

where a_i and b_i are introduced in Lemma 8.5.3.

Proof. When c_i^* and c_j^* are idempotents, the proposed map is obviously well-defined and has inverse $\alpha \mapsto b_i \alpha(c_i^*) a_j$. Now assume that $i = 0$ and $(c_0^*)^2 = 0$, but that c_j^* is an idempotent. By Lemma 9.1.6, it suffices to prove that

$$\phi : \mathrm{Hom}_{\mathcal{A}}(j, 0) \rightarrow c_0^* A c_j^*; \quad x \mapsto a_0 x b_j,$$

is an isomorphism. An inverse to ϕ is constructed by mapping any diagram d in $c_0^* A c_j^*$ to $d' a_j$ where d' is the diagram obtained from d by omitting the top row (essentially forgetting the diagram a_0), proving that ϕ is an isomorphism. The case where c_i^* is an idempotent, but c_j^* is not follows similarly. Finally, if $i = j = 0$ and $(c_0^*)^2 = 0$, the statement follows from Lemma 9.1.6 if the following is an isomorphism,

$$\mathrm{Hom}_{\mathcal{A}}(0, 0) \rightarrow c_0^* A \cap A c_0^*; \quad x \mapsto a_0 x b_0.$$

Both spaces are one-dimensional and the above just maps the empty diagram in $\mathrm{Hom}_{\mathcal{A}}(0, 0)$ to $c_0^* = a_0 b_0$, showing that also here we obtain an isomorphism. $\square$

Proof of Theorem 9.1.1. Consider the isomorphism in Lemma 9.1.7 and take $x \in \mathrm{Hom}_{\mathcal{A}}(j, i)$ and $y \in \mathrm{Hom}_{\mathcal{A}}(k, j)$, so $xy \in \mathrm{Hom}_{\mathcal{A}}(k, i)$. We claim that

$$\alpha_y \circ \alpha_x = \alpha_{xy}.$$

We have $\alpha_y \circ \alpha_x(c_i^*) = \alpha_y(a_i x b_j)$. We can take $x' \in \mathrm{Hom}_{\mathcal{A}}(n, i)$ such that $x' a_j = x$, then

$$\alpha_y(a_i x b_j) = a_i x' \alpha_y(c_j^*) = a_i x' a_j y b_k = a_i x y b_k,$$

proving the claim. $\square$

9.2. Covers.

Theorem 9.2.1. *Let C be any of the category algebras in Lemma 8.5.3 and $A = e_n^* C e_n^*$ the corresponding diagram algebra. Then C is a cover of A .*

Proof. The condition in case (a) of Section 2.4 follows from Corollary 9.1.2. $\square$

In most cases, the cover C is Morita equivalent to A , by Theorem 8.5.1. Now we focus on the remaining cases. Consider the full subcategory $\overline{\mathcal{F}}$ of $C\text{-mod}$ of modules which admit a filtration with sections $\overline{\Delta}(\mu)$, $\mu \in \Lambda$. Recall from Section 2.4 the exact functor

$$F := \mathrm{Hom}_C(Ce_n^*, -) \cong e_n^* - \cong e_n^* C \otimes_C - : C\text{-mod} \rightarrow A\text{-mod}.$$

Theorem 9.2.2. *Let A be $\mathcal{P}_n(0)$ with $n > 2$; $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$ with n even and $n > 4$; or $A = \mathcal{B}_{r,r}(0)$ with $r > 2$. For all $M, N \in \overline{\mathcal{F}}$, the functor F induces isomorphisms*

$$\text{Hom}_C(M, N) \xrightarrow{\sim} \text{Hom}_A(FM, FN) \quad \text{and} \quad \text{Ext}_C^1(M, N) \xrightarrow{\sim} \text{Ext}_A^1(FM, FN).$$

Hence, the cover C is 1-faithful.

We make preparations for the proof of the theorem. Recall, from Section 2.4, that the left exact functor

$$G' := \text{Hom}_A(e_n^*C, -) : A\text{-mod} \rightarrow C\text{-mod},$$

is right adjoint to F . Hence we have the adjoint (unit) natural transformation

$$\eta : \text{Id} \rightarrow G' \circ F.$$

Lemma 9.2.3. *Let A be $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$, for n even, $\mathcal{P}_n(0)$ or $\mathcal{B}_{r,r}(0)$.*

(1) *For any M in $C\text{-mod}$, η_M induces an isomorphism $M \cong G' \circ F(M)$ if and only if*

$$e_0^*M \rightarrow \text{Hom}_A(Ac_0^*, e_n^*M) : \quad v \mapsto \beta_v, \quad \text{where } \beta_v(c_0^*) = a_0v,$$

is an isomorphism, with a_0 from Lemma 8.5.3.

(2) *For any M in $C\text{-mod}$, we have $\mathcal{R}_1 G' \circ F(M) = 0$ if and only if*

$$\text{Ext}_A^1(Ac_0^*, e_n^*M) = 0.$$

Proof. We evaluate η on M , to get the following morphism of C -modules:

$$\eta_M : M \rightarrow G' \circ F(M) = \text{Hom}_A(e_n^*C, e_n^*M); \quad v \mapsto \alpha_v, \quad \alpha_v(c) = cv.$$

This restricts to vector space morphisms

$$\eta^i : e_i^*M \rightarrow \text{Hom}_A(e_n^*C e_i^*, e_n^*M) \quad \text{for } i \in I_A.$$

We also introduce

$$\rho^i : \text{Hom}_A(e_n^*C e_i^*, e_n^*M) \rightarrow e_i^*M; \quad \alpha \mapsto b_i \alpha(a_i),$$

with a_i, b_i as in Lemma 8.5.3. Then η^i and ρ^i are mutually inverse unless $i = 0$.

Thus we find that η_M is an isomorphism if and only if η^0 is. Part (1) then follows from applying the isomorphism $\text{Hom}_A(e_n^*C e_0^*, e_n^*M) \cong \text{Hom}_A(Ac_0^*, e_n^*M)$ from Corollary 9.1.2.

To prove part (2), we observe that we have

$$\mathcal{R}_1 G' \cong \text{Ext}_A^1(e_n^*C, -) \cong \text{Ext}_A^1\left(\bigoplus_{j \in I_A} Ac_j^*, -\right) \cong \text{Ext}_A^1(Ac_0^*, -),$$

by Corollary 9.1.2, since Ac_j^* is projective for $j \neq 0$. □

The proof of the lemma also gives the following result.

Corollary 9.2.4. *Let A be as in Lemma 9.2.3. For $i \neq 0$, we have*

$$\text{Hom}_A(Ac_i^*, e_n^*M) \cong e_i^*M, \quad \text{for any } M \in A\text{-mod}.$$

9.2.5. We fix some of the c_i^* in Lemma 8.5.3. For $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$ with n even, take

$$\begin{aligned} c_0^* &= \begin{array}{ccccccc} \cup & \cup & \cup & \cdots & \cup \\ \cap & \cap & \cap & \cdots & \cap \end{array} & c_2^* &= \begin{array}{c} | \quad \cup \quad \cup \quad \cdots \quad \cup \\ \cap \quad \cap \quad \cdots \quad \cap \end{array} \\ c_4^* &= \begin{array}{c} | \quad | \quad | \quad \cup \quad \cdots \quad \cup \\ \cap \quad \cap \quad \cdots \quad \cap \end{array} . \end{aligned}$$

For $A = \mathcal{B}_{r,r}(0)$, take

$$\begin{aligned} c_0^* &:= \begin{array}{c} \cdots \quad \cup \quad \cdots \\ \cap \quad \cap \quad \cap \quad \cap \quad \cap \\ \cdots \quad \cup \quad \cdots \end{array} & c_1^* &:= \begin{array}{c} \cdots \quad \cup \quad \cdots \\ \cap \quad \cap \quad \cap \quad \cap \quad \cap \\ \cdots \quad \cup \quad \cdots \end{array} \\ c_2^* &:= \begin{array}{c} | \quad \cdots \quad \cup \quad \cdots \\ \cap \quad \cap \quad \cap \quad \cap \quad \cap \\ | \quad \cdots \quad \cup \quad \cdots \end{array} . \end{aligned}$$

For $\mathcal{P}_n(0)$, take

$$\begin{aligned} c_0^* &= \{\{1, 2, \dots, n\}, \{1', 2', \dots, n'\}\}, & c_1^* &= \{\{1, 1', 2', \dots, n'\}, \{2, 3, \dots, n\}\}, \text{ and} \\ c_2^* &= \{\{1, 1'\}, \{2, 2', 3', \dots, n'\}, \{3, 4, \dots, n\}\}. \end{aligned}$$

We also introduce $\gamma = \gamma_A$, where $\gamma_A = 1$ for $A = \mathcal{B}_{r,r}(0)$ or $\mathcal{P}_n(0)$ and $\gamma_A = 2$ otherwise.

Lemma 9.2.6. *Consider A equal to $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$ for n even and $n \geq 4$, $\mathcal{P}_n(0)$ with $n \geq 2$, or $\mathcal{B}_{r,r}(0)$ with $r \geq 2$. We have*

$$\text{Hom}_A(Ac_0^*, A/Ac_0^*A) = 0.$$

Proof. This is a stronger version of Lemma 9.1.6. It can be proved using the same arguments. Consider $\alpha : Ac_0^* \rightarrow A/Ac_0^*A$ and $a \in A$ such that $a + Ac_0^*A = \alpha(c_0^*)$.

For $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$, we have $c_2^*c_0^* = c_0^*$, so we can assume that $a \in c_2^*A$. As w in equation (9.4) is an idempotent satisfying $wc_0^* = c_0^*$, we can further assume that $wa = a$. These two conditions on a show that it must be a linear combination of diagrams with $n/2$ caps, so $a \in Ac_0^*A$ and hence $\alpha = 0$.

The proof for $\mathcal{B}_{r,r}(0)$ is identical, by using the idempotent v in equation (9.5). Also the proof for $\mathcal{P}_n(0)$ works along the same lines, by using $uc_0^* = 0$ with $u = u_1 - u_2$ in (9.2). $\square$

Lemma 9.2.7. *Let A be as in Lemma 9.2.6. There exists an exact sequence*

$$Ac_{2\gamma}^* \rightarrow Ac_\gamma^* \rightarrow Ac_0^* \rightarrow 0.$$

Proof. First, let A be $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$. The map $Ac_2^* \twoheadrightarrow Ac_0^*$ is defined as $a \mapsto ac_0^*$ for any $a \in Ac_2^*$, where surjectivity follows from $c_2^*c_0^* = c_0^*$. Now we determine the kernel K of this epimorphism. From the structure of c_2^* and c_0^* it follows that K is the spanned by all diagrams without propagating lines and by all elements of the form $d_1 - d_2$, where $d_1, d_2 \in Ac_2^*$ are diagrams satisfying the following conditions. For $k \in \{1, 2\}$, the diagram d_k has two propagating lines, connecting 1 to i_k and n to j_k , giving four different dots $\{i_1, i_2, j_1, j_2\}$. There is a cap in d_k which connects dots i_l and j_l , with $\{k, l\} = \{1, 2\}$. Finally, removing these caps and propagating lines in d_1 and d_2 yield identical $(n-2, n-4)$ -Brauer diagrams.

An example of such a $d_1 - d_2$, with $i_1 = 1'$, $j_1 = 2'$, $i_2 = 3'$ and $j_2 = 4'$, is given by

$$x := \left| \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \right| - \left| \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} \right|$$

The diagrams consist of horizontal lines with various arcs (cups and caps) and vertical propagating lines. Diagram 1 and 2 are connected by a vertical line on the left. Diagram 3 and 4 are connected by a vertical line on the left. The minus sign indicates the difference between the two pairs of diagrams.

Now we claim that $K = Ax$. That the span of all diagrams without propagating lines is in Ax follows easily from multiplying x with the diagram having a cup and cap connecting the first two dots and otherwise only vertical propagating lines. For $d_1 - d_2 \in K$ as above, we consider the three algebras separately.

For $A = \mathcal{B}_n(0)$, we can consider a diagram $a \in \mathbb{S}_n$, where 1 is connected to i_1 , 2 to j_1 , 3 to i_2 and 4 to j_2 . It then follows easily that a can be completed such that $ax = d_1 - d_2$.

For $A = \text{TL}_n(0)$, take an arbitrary $d_1 - d_2$ as above and consider the $n/2 - 2$ cups which appear in both diagrams d_1 and d_2 . It follows easily that this information determines $d_1 - d_2$ uniquely, up to sign. Now we consider the unique diagram $a \in A$ which contains those $n/2 - 2$ cups, the $n/2 - 2$ caps which appear in c_4^* and four propagating lines. Then we find $ax = \pm(d_1 - d_2)$. Finally, the case $A = J_n(0)$ follows similarly.

As $c_4^*x = x$, we have a surjection $Ac_4^* \twoheadrightarrow K$, with $K = Ax$, proving the exact sequence.

For the two remaining algebras one proves, similarly to the above, that the kernel of $Ac_1^* \twoheadrightarrow Ac_0^*$ is generated by $x = c_2^*x$, given by

$$x := \left| \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} \right| - \left| \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} \right|$$

The diagrams show a central vertical line with various arcs and other lines crossing it. Diagram 5 and 6 are connected by a vertical line on the left. Diagram 7 and 8 are connected by a vertical line on the left. The minus sign indicates the difference between the two pairs of diagrams.

for $A = \mathcal{B}_{r,r}(0)$, and by

$$x := \{\{1, 1'\}, \{2, 3, \dots, n\}, \{2', 3', \dots, n'\}\} - \{\{1'\}, \{2, 3, \dots, n\}, \{1, 2', 3', \dots, n'\}\},$$

for $A = \mathcal{P}_n(0)$. □

Lemma 9.2.8. *For A as in Lemma 9.2.3, let $D_i := e_n^*C \otimes_B He_i^*$. Then*

$$D_i \cong Ac_i^*/Ac_{i-\gamma}^*Ac_i^*, \quad \text{for } i \in I_A.$$

Proof. Set $f' = \sum_{j < i} e_j^*$ and $f = f' + e_i^*$. Lemma 3.2.8 then implies that

$$C \otimes_B He_i^* \cong (C/Cf'C)f.$$

Corollary 9.1.2 and Theorem 9.1.1 then imply the claim. □

Proposition 9.2.9. *For A in Lemma 9.2.6, the unit $\eta : \text{Id} \rightarrow G' \circ F$ induces isomorphisms*

$$G' \circ F(\overline{\Delta}(i, \nu)) \cong \overline{\Delta}(i, \nu) \quad \text{for all } i \in I_A \text{ and } \nu \in \Lambda_i.$$

Proof. The proof is similar for all algebras. Let A be $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$. For $i = 0$, Lemma 9.2.8 implies that

$$F(\overline{\Delta}(0, \emptyset)) \cong e_n^*Ce_0^* \cong Ac_0^*.$$

By Lemma 9.2.3(1), it thus suffices to prove that

$$e_0^*Ce_0^* \rightarrow \text{Hom}_A(Ac_0^*, Ac_0^*) : \quad v \mapsto \beta_v, \quad \text{where } \beta_v(c_0^*) = a_0vb_0,$$

is an isomorphism. This is Lemma 9.1.7 for $i = j = 0$.

For $i > 0$, we have $e_0^*\overline{\Delta}(i, \nu) = 0$, so by Lemma 9.2.3(1), we only need to prove

$$(9.6) \quad \text{Hom}_A(Ac_0^*, e_n^*\overline{\Delta}(i, \nu)) = 0.$$

Recall that He_2^* is a group algebra and hence self-injective. In particular, for any simple module $L^0(2, \nu)$ of He_2^* we have $L^0(2, \nu) \hookrightarrow He_2^*$ and consequently $e_n^* \overline{\Delta}(2, \nu) \hookrightarrow D_2$. By Lemma 9.2.8, D_2 is a direct summand of A/Ac_0^*A . Hence Lemma 9.2.6 implies

$$\text{Hom}_A(Ac_0^*, D_2) = 0.$$

Thus the above equation implies equation (9.6), concluding the case $i = 2$. Finally, for $i > 2$, we use Lemma 9.2.7, which implies that the left-hand side of equation (9.6) is a subspace of $\text{Hom}_A(Ac_2^*, e_n^* \overline{\Delta}(i, \nu))$, which is equal to $e_2^* \overline{\Delta}(i, \nu)$ by Corollary 9.2.4. Now $e_2^* \overline{\Delta}(i, \nu) = 0$ since $i > 2$, and equation (9.6) is again satisfied. $\square$

9.2.10. From now on we will assume that for A equal to $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$ we have $n > 4$. This allows to introduce diagrams $y_1, y_2 \in A$ as

$$y_1 := \left| \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \right| \quad y_2 := \left| \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} \right|$$

Similarly, for $A = \mathcal{B}_{r,r}(0)$ we will assume $r > 2$, which allows to introduce

$$y_1 := \left| \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} \right| \quad y_2 := \left| \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} \right|$$

For all four algebras a direct computation shows that

$$(9.7) \quad x = y_1 x - y_2 x,$$

with x as introduced in the proof of Lemma 9.2.7.

For $A = \mathcal{P}_n(0)$ we will assume that $n > 2$ and we introduce $y \in A$,

$$y := \{ \{1, 1'\}, \{2, 2'\}, \{3, 4, \dots, n\}, \{3', 4', \dots, n'\} \} - \{ \{1, 1'\}, \{2'\}, \{3, 4, \dots, n\}, \{2, 3', 4', \dots, n'\} \} + \{ \{1'\}, \{1, 2'\}, \{3, 4, \dots, n\}, \{2, 3', 4', \dots, n'\} \}.$$

It follows immediately that $yx = 0$.

Lemma 9.2.11. *Let A be as in Theorem 9.2.2. Then*

$$\text{Ext}_A^1(Ac_0^*, Ac_0^*) = 0.$$

Proof. First we will prove $\text{Hom}_A(Ax, Ac_0^*) = 0$. Consider $\phi : Ax \rightarrow Ac_0^*$ and $a := \phi(c_0^*)$.

First, let A be $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$, the case $\mathcal{B}_{r,r}(0)$ is proved similarly. As $c_4^* x = x$, we have $a \in c_4^* Ac_0^*$. By equation (9.7), we must have

$$a = y_1 a - y_2 a.$$

This means that a is a linear combination of diagrams which have the $n/2$ caps of c_0^* , the $n/2 - 2$ cups of c_4^* and another cup connecting either $1', 2'$, or $3', 4'$. The only such diagram is c_0^* . However, $y_1 c_0^* = y_2 c_0^*$ and hence $a = 0$.

For $A = \mathcal{P}_n(0)$, we have $a \in c_2^* Ac_0^*$, with $\dim c_2^* Ac_0^* = 2$. With y from 9.2.10, we must have $ya = 0$. It follows by direct computation that $ya = 0$ for $a \in c_2^* Ac_0^*$ implies $a = 0$.

Hence, in every case we have indeed, $\text{Hom}_A(Ax, Ac_0^*) = 0$. Recall the short exact sequence

$$(9.8) \quad 0 \rightarrow Ax \rightarrow Ac_\gamma^* \rightarrow Ac_0^* \rightarrow 0$$

from the proof of Lemma 9.2.7. This implies an exact sequence

$$(9.9) \quad \text{Hom}_A(Ax, M) \rightarrow \text{Ext}_A^1(Ac_0^*, M) \rightarrow 0,$$

for any A -module M , as Ac_γ^* is projective. As we established that $\text{Hom}_A(Ax, Ac_0^*) = 0$, the statement follows. $\square$

Lemma 9.2.12. *Let A be as in Theorem 9.2.2. Then*

$$\text{Ext}_A^1(Ac_0^*, Ac_{2\gamma}^*/Ac_\gamma^*Ac_{2\gamma}^*) = 0.$$

Proof. We will prove $\text{Hom}_A(Ax, Ac_{2\gamma}^*/Ac_\gamma^*Ac_{2\gamma}^*) = 0$, then the statement follows from (9.9). Consider $\phi : Ax \rightarrow Ac_{2\gamma}^*/Ac_\gamma^*Ac_{2\gamma}^*$ and $a := \phi(c_0^*)$.

First, let A be $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$, the case $\mathcal{B}_{r,r}(0)$ is proved similarly. As $c_4^*x = x$, we have $a \in c_4^*A/(Ac_2^*)c_4^*$. By equation (9.7), we must have

$$a = y_1a - y_2a.$$

This means that a is represented by a linear combination of diagrams, where each contains 4 propagating lines, the $n/2 - 2$ cups and caps of c_4^* , but also either the cup connecting $1', 2'$ or $3', 4'$. This is an inconsistency, so $a = 0$.

For $A = \mathcal{P}_n(0)$, the dimension of $c_2^*(A/Ac_0^*A)c_2^*$ is 2. It follows quickly that no non-zero element a satisfies $ya = 0$, so $a = 0$. $\square$

Lemma 9.2.13. *Let A be as in Theorem 9.2.2. Then*

$$\text{Ext}_A^1(Ac_0^*, Ac_\gamma^*/Ac_0^*Ac_\gamma^*) = 0.$$

Proof. Set $M := Ac_\gamma^*/Ac_0^*Ac_\gamma^*$. The short exact sequence (9.8) and Lemma 9.2.6 imply a short exact sequence

$$0 \rightarrow \text{Hom}_A(Ac_\gamma^*, M) \rightarrow \text{Hom}_A(Ax, M) \rightarrow \text{Ext}_A^1(Ac_0^*, M) \rightarrow 0.$$

We will prove that for each algebra,

$$\dim \text{Hom}_A(Ax, M) \leq d := \dim c_\gamma^*(A/Ac_0^*A)c_\gamma^* = \dim He_\gamma^*,$$

proving that the extension group must vanish.

First, let A be $\mathcal{B}_n(0)$, $J_n(0)$ or $\text{TL}_n(0)$, the case $\mathcal{B}_{r,r}(0)$ is proved similarly. If a is the image of x under a morphism $Ax \rightarrow M$, then, by equation (9.7), we have

$$a = c_4^*a \quad \text{and} \quad a = y_1a - y_2a.$$

Hence a must be represented by a linear combinations of diagrams, containing the $n/2 - 1$ caps of c_2^* , the $n/2 - 2$ cups of c_4^* and a cup which either connects $1', 2'$ or $3', 4'$. The dimension of the space of such a is 4 for $\mathcal{B}_n(0)$ and $J_n(0)$ and 2 for $\text{TL}_n(0)$. In each case, imposing the actual condition that $a = y_1a - y_2a$, leaves half of the dimensions. For each case, this yields precisely d .

For $A = \mathcal{P}_n(0)$, we have $d = 1$ and the dimension of $c_2^*(A/Ac_0^*A)c_1^*$ is 3. The subspace of elements that are annihilated by left multiplication with y also has dimension 1. $\square$

Proposition 9.2.14. *Maintain the notation and assumptions of Theorem 9.2.2. We have*

$$\mathcal{R}_1G' \circ F(\overline{\Delta}(i, \nu)) = 0, \quad \text{for all } i \in I_A \text{ and } \nu \in \Lambda_i.$$

Proof. By Lemma 9.2.3(2), it suffices to prove that

$$(9.10) \quad \text{Ext}_A^1(Ac_0^*, e_n^*\overline{\Delta}(i, \nu)) = 0.$$

By Lemma 9.2.7, this space is a subquotient of

$$\text{Hom}_A(Ac_{2\gamma}^*, e_n^*\overline{\Delta}(i, \nu)),$$

which is zero when $i > 2\gamma$, by Corollary 9.2.4. Hence we focus on $i \in \{0, \gamma, 2\gamma\}$.

By Lemma 9.2.8, we have $e_n^* \overline{\Delta}(0, \emptyset) \cong Ac_0^*$, so (9.10) is satisfied for $i = 0$ by Lemma 9.2.11.

As He_i^* is self-injective, for D_i in Lemma 9.2.8, we have a short exact sequence

$$0 \rightarrow e_n^* \overline{\Delta}(i, \nu) \rightarrow D_i \rightarrow Q_\nu \rightarrow 0,$$

where Q_ν has a filtration with sections $\overline{\Delta}(i, \nu')$ with $\nu' \in \Lambda_i$. This gives an exact sequence

$$\mathrm{Hom}_A(Ac_0^*, Q_\nu) \rightarrow \mathrm{Ext}_A^1(Ac_0^*, e_n^* \overline{\Delta}(i, \nu)) \rightarrow \mathrm{Ext}_A^1(Ac_0^*, D_i).$$

If $i \neq 0$, the left-hand space is zero by (9.6). For $i \in \{\gamma, 2\gamma\}$, the right-hand side is zero by Lemmata 9.2.12 and 9.2.13. Hence the middle term is zero and (9.10) is satisfied. $\square$

Proof of Theorem 9.2.2. By Proposition 9.2.14, we have

$$\mathcal{R}_1 G'(FM) = \mathrm{Ext}_A^1(e_n^* C, FM) = 0, \quad \text{for any } M \in \overline{\mathcal{F}}.$$

Consider a short exact sequence $M_1 \hookrightarrow M \twoheadrightarrow M_2$, with M_i (and hence also M) in $\overline{\mathcal{F}}$. Using the above vanishing of cohomology, we find a commutative diagram with exact rows

$$(9.11) \quad \begin{array}{ccccccc} 0 & \longrightarrow & M_1 & \longrightarrow & M & \longrightarrow & M_2 \longrightarrow 0 \\ & & \downarrow \eta_{M_1} & & \downarrow \eta_M & & \downarrow \eta_{M_2} \\ 0 & \longrightarrow & G'FM_1 & \longrightarrow & G'FM & \longrightarrow & G'FM_2 \longrightarrow 0. \end{array}$$

This implies that, if η_{M_1} and η_{M_2} are isomorphisms, so is η_M . Proposition 9.2.9 can then be used to prove, by induction on the length of the filtration that we have

$$\eta_M : M \xrightarrow{\sim} G' \circ F(M),$$

for any $M \in \overline{\mathcal{F}}$. Hence, we have an isomorphism

$$\mathrm{Hom}_C(M, N) \cong \mathrm{Hom}_C(M, G' \circ F(N)) \cong \mathrm{Hom}_A(FM, FN),$$

induced by F . Now we consider arbitrary $M_1, M_2 \in \overline{\mathcal{F}}$. The morphism

$$F : \mathrm{Ext}_C^1(M_2, M_1) \rightarrow \mathrm{Ext}_A^1(FM_2, FM_1)$$

has left inverse induced by G' , by (9.11). As furthermore $F \circ G' \cong \mathrm{Id}$, the above morphism is an isomorphism. This concludes the proof. $\square$

9.3. Cell modules and standard systems.

Theorem 9.3.1. *Assume that the field $\mathbb{k}$ is algebraically closed. Let A be $\mathcal{B}_n(0)$, $J_n(0)$ or $\mathrm{TL}_n(0)$ with n even, $\mathcal{P}_n(0)$ or $\mathcal{B}_{r,r}(0)$. The cell modules of the cellular algebra A form a standard system if and only if the following condition is satisfied.*

algebra A	condition
$\mathcal{P}_n(0)$	$n > 2$ and $\mathrm{char}(\mathbb{k}) \notin \{2, 3\}$
$\mathcal{B}_n(0)$	$n \notin \{2, 4\}$ and $\mathrm{char}(\mathbb{k}) \notin \{2, 3\}$
$J_n(0)$	$n \notin \{2, 4\}$ and $\mathrm{char}(\mathbb{k}) \notin [2, n/2]$
$\mathrm{TL}_n(0)$	$n \notin \{2, 4\}$
$\mathcal{B}_{r,r}(0)$	$r > 2$ and $\mathrm{char}(\mathbb{k}) \notin \{2, 3\}$

First we prove the following lemma.

Lemma 9.3.2. *The cell modules of*

- (1) $\mathcal{B}_2(0)$, $J_2(0)$, $\mathrm{TL}_2(0)$ and $\mathcal{B}_{1,1}(0)$ do not form a standard system, for any partial order;

(2) $\mathcal{B}_4(0)$, $J_4(0)$, $\mathrm{TL}_4(0)$, $\mathcal{P}_2(0)$ and $\mathcal{B}_{2,2}(0)$ do not form a standard system.

Proof. First we prove part (1), we have

$$\mathrm{TL}_2(0) \cong \mathcal{B}_{1,1}(0) \cong \mathbb{k}[x](x^2).$$

The only cellular structure on this algebra gives two cell modules which are isomorphic. If $\mathrm{char}(\mathbb{k}) \neq 2$, we have

$$\mathcal{B}_2(0) \cong J_2(0) \cong \mathbb{k}[x]/(x^2) \oplus \mathbb{k},$$

so the result follows as above. If $\mathrm{char}(\mathbb{k}) = 2$, the algebra $\mathcal{B}_2(0) \cong J_2(0)$ has two isomorphic cell modules, induced from the sign and trivial $\mathbb{k}\mathbb{S}_2$ -module, by Proposition 8.6.4.

Now we prove part (2). We have $c_{2\gamma}^* = 1$ and, as c_γ^* is an idempotent, equation (9.8) implies

$$\mathrm{Hom}_A(Ax, A/Ac_\gamma^*A) \cong \mathrm{Ext}^1(Ac_0^*, A/Ac_\gamma^*A).$$

As $Ac_0^* \cong W(0, \emptyset)$ and A/Ac_γ^*A has a filtration with sections of the form $W(2\gamma, \nu)$ with $\nu \in L_{2\gamma}$, the right-hand side must be zero in order to have a standard system for $(L, \leq)$. However, we claim that there exists a non-zero morphism

$$\phi : Ax \rightarrow A/Ac_\gamma^*A.$$

For $A = \mathrm{TL}_4(0)$, we can set $\phi(x) = 1$. For the other cases, x is mapped to an appropriate anti-symmetrisation of 1. $\square$

Proof of Theorem 9.3.1. By proposition 8.6.4, under the conditions in Theorem 9.2.2, the cell modules of A form a standard system if and only if the cell modules of C form a standard system. The necessary and sufficient condition for the latter is given in Theorem 8.4.3.

For the remaining cases, *i.e.* when Theorem 9.2.2 is not applicable, the cell modules do not form a standard system by Lemma 9.3.2. $\square$

APPENDIX A. STRATIFIED ALGEBRAS

A.1. Homological stratification. An *exact stratification of an algebra A* is a chain (2.4) or ideals such that J_i/J_{i-1} is exactly stratifying in A/J_{i-1} , see 2.6.1(0), for $1 \leq i \leq m$. For a chain (2.4) we consider the algebras $A^{(i)}$ as in Remark 2.6.5 and we identify $A^{(i)}$ -modules with $f_{i+1}A f_{i+1}$ -modules having trivial $f_{i+1}J_i f_{i+1}$ -action. The following principle is well-known.

Lemma A.1.1. *Consider an algebra A with an exact stratification (2.4). For any $A^{(l)}$ -module M and $A^{(j)}$ -module N , with $0 \leq l \leq j \leq m-1$, let $\widetilde{M} := A f_{l+1} \otimes_{f_{l+1}A f_{l+1}} M$ and $\widetilde{N} := A f_{j+1} \otimes_{f_{j+1}A f_{j+1}} N$. Then*

$$\mathrm{Ext}_A^k(\widetilde{M}, \widetilde{N}) \cong \delta_{jl} \mathrm{Ext}_{A^{(l)}}^k(M, N), \quad \text{for any } k \in \mathbb{N}.$$

Proof. First we claim that $J_l \widetilde{M} = 0 = J_l \widetilde{N}$. It suffices to show that f_l acts as zero. Now f_l is included in $f_{j+1}J_l f_{j+1}$ since $l \leq j$. Action of $f_{j+1}J_l f_{j+1}$ on $\widetilde{N}$ gives

$$f_{j+1}J_l f_{j+1} \otimes_{f_{j+1}A f_{j+1}} N = 0.$$

This observation and [CPS2, equation (2.1.2.1)] imply

$$\mathrm{Ext}_A^k(\widetilde{M}, \widetilde{N}) \cong \mathrm{Ext}_{A/J_l}^k(\widetilde{M}, \widetilde{N}).$$

Now, as A/J_l -modules, we have

$$\widetilde{M} \cong (A/J_l)f_l \otimes_{A^{(l)}} M \quad \text{and} \quad \widetilde{N} \cong (A/J_l)f_{j+1} \otimes_{f_{j+1}(A/J_l)f_{j+1}} N.$$

As the right $A^{(l)}$ -module $(A/J_l)f_{l+1}$ is projective by Lemma A.1.2(2), we can use equation (2.2) to obtain

$$\mathrm{Ext}_{A/J_l}^k(\widetilde{M}, \widetilde{N}) \cong \mathrm{Ext}_{A^{(l)}}^k(M, f_{l+1}(A/J_l)f_{j+1} \otimes_{f_{j+1}(A/J_l)f_{j+1}} N).$$

As we have

$$f_{l+1}(A/J_l)f_{j+1} \otimes_{f_{j+1}(A/J_l)f_{j+1}} N \cong \begin{cases} 0 & \text{for } l < j, \\ N & \text{for } l = j, \end{cases}$$

this concludes the proof. $\square$

Lemma A.1.2. *Consider an idempotent ideal $J = AeA$ in A .*

- (1) *The ideal J is standardly stratifying if and only if*
 - *multiplication induces an A -bimodule isomorphism $Ae \otimes_{eAe} eA \xrightarrow{\sim} J$, and*
 - *the left eAe -module eA is projective.*
- (2) *The ideal J is exactly stratifying if and only if*
 - *multiplication induces an A -bimodule isomorphism $Ae \otimes_{eAe} eA \xrightarrow{\sim} J$, and*
 - *the right eAe -module Ae is projective.*

Proof. We prove part (1) as part (2) then follows from considering A^{op} . If $Ae \otimes_{eAe} eA \xrightarrow{\sim} J$, then the left A -module J is induced from the left eAe -module eA . If the latter is projective then ${}_AJ$ is also projective, so the ideal J is standardly stratifying. If J is standardly stratifying, the conclusion follows from the proof of [CPS2, Remark 2.1.2(b)]. $\square$

A.2. Comparison with Kleshchev's terminology. For any class $\mathcal{B}$ of Noetherian, positively graded, connected algebras, Kleshchev introduced in [Kl, Section 6] the notions of $\mathcal{B}$ -standardly stratified, $\mathcal{B}$ -properly stratified and $\mathcal{B}$ -quasi-hereditary algebras. Even though the aim of [Kl] is to study infinite dimensional and graded algebras, the definitions also cover all the cases in Definition 2.7.1. Therefore, we assume that every grading is reduced to the zero component and introduce the classes $\mathcal{S} \subset \mathcal{L} \subset \mathcal{D}$, where $\mathcal{D}$ contains the finite dimensional unital algebras, $\mathcal{L}$ the quasi-local algebras and $\mathcal{S}$ the semisimple algebras. Then we have the following identification between our notions and those in [Kl]:

	$\mathcal{B} = \mathcal{D}$	$\mathcal{B} = \mathcal{L}$	$\mathcal{B} = \mathcal{S}$
$\mathcal{B}$ -standardly stratified	standardly stratified	strongly standardly stratified	quasi-hereditary
$\mathcal{B}$ -properly stratified	exactly standardly stratified	properly stratified	quasi-hereditary
$\mathcal{B}$ -quasi-hereditary	properly stratified	properly stratified	quasi-hereditary

This follows by comparing [Kl, Definitions 6.1 and 6.2] with 2.6.1, using Lemma A.1.2.

Acknowledgement. The research was supported by Australian Research Council Discovery Project Grant DP140103239 and an FWO postdoctoral grant.

The authors thank Steffen König, Julian Külshammer and Volodymyr Mazorchuk for very useful comments on the first version of the manuscript.

REFERENCES

- [ADL] I. Ágoston, V. Dlab, E. Lukács: Stratified algebras. *C. R. Math. Acad. Sci. Soc. R. Can.* **20** (1998), no. 1, 22–28.
- [AST] H.H. Andersen, C. Stroppel, D. Tubbenhauer: Cellular structures using U_q -tilting modules. [arXiv:1503.00224](https://arxiv.org/abs/1503.00224).
- [Au] M. Auslander: Representation theory of Artin algebras. I, II. *Comm. Algebra* **1** (1974), 177–268; *ibid.* **1** (1974), 269–310.
- [APT] M. Auslander, M.I. Platzeck, G. Todorov: Homological theory of idempotent ideals. *Trans. Amer. Math. Soc.* **332** (1992), no. 2, 667–692.
- [BFK] J. Bernstein, I. Frenkel, M. Khovanov: A categorification of the Temperley-Lieb algebra and Schur quotients of $U(\mathfrak{sl}_2)$ via projective and Zuckerman functors. *Selecta Math. (N.S.)* **5** (1999), no. 2, 199–241.
- [Bo] C. Bowman: Bases of quasi-hereditary covers of diagram algebras. *Math. Proc. Cambridge Philos. Soc.* **154** (2013), no. 3, 393–418.
- [Ca] Y. Cao: On the quasi-heredity and the semi-simplicity of cellular algebras. *J. Alg.* **267** (2003) 323–341.
- [CPS1] E. Cline, B. Parshall, L. Scott: Finite-dimensional algebras and highest weight categories. *J. Reine Angew. Math.* **391** (1988), 85–99.
- [CPS2] E. Cline, B. Parshall, L. Scott: Stratifying endomorphism algebras. *Mem. Amer. Math. Soc.* **124** (1996), no. 591.
- [CW] J. Comes, B. Wilson: Deligne’s category $\underline{\mathrm{Rep}}(GL_\delta)$ and representations of general linear supergroups. *Represent. Theory* **16** (2012), 568–609.
- [CM] K. Coulembier, V. Mazorchuk: Some homological properties of category $\mathcal{O}$. III. *Adv. Math.* **283** (2015), 204–231.
- [CDDM] A.G. Cox, M. De Visscher, S. Doty, P.P. Martin: On the blocks of the walled Brauer algebra, *J. Algebra* **320** (2008), 169–212.
- [De] P. Deligne: La catégorie des représentations du groupe symétrique S_t , lorsque t n’est pas un entier naturel. *Algebraic groups and homogeneous spaces*, 209–273, *Tata Inst. Fund. Res. Stud. Math.*, Tata Inst. Fund. Res., Mumbai, 2007.
- [DJM] R. Dipper, G. James, A. Mathas: The (Q, q) -Schur algebra. *Proc. London Math. Soc. (3)* **77** (1998), no. 2, 327–361.
- [Dl] V. Dlab: Properly stratified algebras. *C. R. Acad. Sci. Paris Sér. I Math.* **331** (2000), no. 3, 191–196.
- [DR1] V. Dlab, C.M. Ringel: Every semiprimary ring is the endomorphism ring of a projective module over a quasihereditary ring. *Proc. Amer. Math. Soc.* **107** (1989), no. 1, 1–5.
- [DR2] V. Dlab, C.M. Ringel: The module theoretical approach to quasi-hereditary algebras. *Representations of algebras and related topics (Kyoto, 1990)*, 200–224, *London Math. Soc. Lecture Note Ser.*, **168**, Cambridge Univ. Press, Cambridge, 1992.
- [DuL] J. Du, Z. Lin: Near-matrix filtrations and quasi-hereditary algebras. Unpublished manuscript.
- [DuR] J. Du, H. Rui: Based algebras and standard bases for quasi-hereditary algebras. *Trans. Amer. Math. Soc.* **350** (1998), no. 8, 3207–3235.
- [DuS] J. Du, L. Scott: The q -Schur² algebra. *Trans. Amer. Math. Soc.* **352** (2000), no. 9, 4325–4353.
- [Fr] A. Frisk: Dlab’s theorem and tilting modules for stratified algebras. *J. Algebra* **314** (2007), no. 2, 507–537.
- [GL] J.J. Graham, G.I. Lehrer: Cellular algebras. *Invent. Math.* **123** (1996), no. 1, 1–34.
- [HHKP] R. Hartmann, A. Henke, S. König, R. Paget: Cohomological stratification of diagram algebras. *Math. Ann.* **347** (2010), no. 4, 765–804.
- [HP] R. Hartmann, R. Paget: Young modules and filtration multiplicities for Brauer algebras. *Math. Z.* **254** (2006), no. 2, 333–357.
- [HN] D.J. Hemmer, D. Nakano: Specht filtrations for Hecke algebras of type A. *J. London Math. Soc. (2)* **69** (2004), no. 3, 623–638.
- [HK] A. Henke, S. König: Schur algebras of Brauer algebras I. *Math. Z.* **272** (2012), no. 3-4, 729–759.

- [Hu] J.E. Humphreys: Representations of semisimple Lie algebras in the BGG category $\mathcal{O}$. Graduate Studies in Mathematics, **94**. American Mathematical Society, Providence, RI, 2008.
- [Ja] G. James: The representation theory of the symmetric groups. Springer LNM **682**, 1978.
- [Jo] V.F.R. Jones: The Potts model and the symmetric group. Subfactors (Kyuzeso, 1993), 259–267, World Sci. Publ., River Edge, NJ, 1994.
- [KKM] O. Khomenko, S. König, V. Mazorchuk: Finitistic dimension and tilting modules for stratified algebras. J. Algebra **286** (2005), no. 2, 456–475.
- [Kl] A.S. Kleshchev: Affine highest weight categories and affine quasihereditary algebras. Proc. Lond. Math. Soc. (3) **110** (2015), no. 4, 841–882.
- [KN] A. Kleshchev, D. Nakano: On comparing the cohomology of general linear and symmetric groups. Pacific J. Math. **201** (2001), no. 2, 339–355.
- [KM] M. Klucznik, V. Mazorchuk: Parabolic decomposition for properly stratified algebras. J. Pure Appl. Algebra **171** (2002), no. 1, 41–58.
- [Kö] S. König: Exact Borel subalgebras of quasi-hereditary algebras. I. With an appendix by Leonard Scott. Math. Z. **220** (1995), no. 3, 399–426.
- [KKO] S. König, J. Külshammer, S. Ovsienko: Quasi-hereditary algebras, exact Borel subalgebras, A_∞ -categories and boxes. Adv. Math. **262** (2014), 546–592.
- [KX1] S. König, C.C. Xi: On the structure of cellular algebras. Algebras and modules, II (Geiranger, 1996), 365–386, CMS Conf. Proc., **24**, Amer. Math. Soc., Providence, RI, 1998.
- [KX2] S. König, C.C. Xi: When is a cellular algebra quasi-hereditary? Math. Ann. **315** (1999), no. 2, 281–293.
- [La] T.Y. Lam: A first course in noncommutative rings. Second edition. Graduate Texts in Mathematics, **131**. Springer-Verlag, New York, 2001.
- [LZ] G.I. Lehrer, R.B. Zhang: The Brauer category and invariant theory. J. Eur. Math. Soc. **17** (2015), 2311–2351.
- [Ma1] P.P. Martin: Temperley-Lieb algebras for nonplanar statistical mechanics - the partition algebra construction. J. Knot Theory Ramifications **3** (1994), no. 1, 51–82.
- [Ma2] P.P. Martin: The structure of the partition algebras. J. Algebra **183** (1996), 319–358.
- [PSW] B. Parshall, L. Scott, J.P. Wang: Borel subalgebras redux with examples from algebraic and quantum groups. Algebr. Represent. Theory **3** (2000), no. 3, 213–257.
- [Ro] R. Rouquier: q -Schur algebras and complex reflection groups. Mosc. Math. J. **8** (2008), no. 1, 119–158, 184.
- [Ru] H. Rui: A criterion on the semisimple Brauer algebras. J. Combin. Theory Ser. A **111** (2005), no. 1, 78–88.
- [So] W. Soergel: Équivalences de certaines catégories de $\mathfrak{g}$ -modules. C. R. Acad. Sci. Paris Sér. I Math. **303** (1986), no. 15, 725–728.
- [We] C.A. Weibel: An introduction to homological algebra. Cambridge Studies in Advanced Mathematics, **38**. Cambridge University Press, Cambridge, 1994.
- [Xi] C.C. Xi: Partition algebras are cellular. Comp. Math. **119** (1999), no. 1, 99–109.
- [YL] G. Yang, Y. Li: Standardly based algebras and 0-Hecke algebras. J. Algebra Appl. **14** (2015), no. 10, 1550141, 13 pp.

K. Coulembier `kevin.coulembier@sydney.edu.au`

School of Mathematics and Statistics, University of Sydney, NSW 2006, Australia

R. B. Zhang `ruibin.zhang@sydney.edu.au`

School of Mathematics and Statistics, University of Sydney, NSW 2006, Australia